

#### www.immunoway.com

# Crk II Mouse mAb

CatalogNo: YM0167

### Key Features

Host Species

Mouse

Reactivity

Human

Applications • WB,IHC,IF,FC,ELISA

MW • 34kD (Calculated)

### **Recommended Dilution Ratios**

WB 1:500-1:2000 IHC 1:200-1:1000 IF 1:200-1:1000 Flow Cyt 1:200-1:400 ELISA 1:10000 Not yet tested in other applications.

### **Storage**

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### **Basic Information**

Clonality Monoclonal

#### Immunogen Information

ImmunogenPurified recombinant fragment of human Crk II expressed in E. Coli.SpecificityCrk II Monoclonal Antibody detects endogenous levels of Crk II protein.

# Target Information

Gene name	CRK		
Protein Name	Adapter molecule crk Organism	Gene ID	UniProt ID
	Human	<u>1398;</u>	<u>P46108;</u>
	Mouse	12928;	<u>Q64010;</u>
Cellular Localization	Cytoplasm . Cell membrane . Translocated to the plasma membrane upon cell adhesion		
Tissue specificity	Embryonic lung,Epithelium,Eye,Lung,Placenta,		
Function	Domain:The C-terminal SH3 domain function as a negative modulator for transformation and the N-terminal SH3 domain appears to function as a positive regulator for transformation.,Domain:The SH2 domain mediates interaction with SHB.,Function:The Crk-I and Crk-II forms differ in their biological activities. Crk-II has less transforming activity than Crk-I. Crk-II mediates attachment-induced MAPK8 activation, membrane ruffling and cell motility in a Rac-dependent manner. Involved in phagocytosis of apoptotic cells and cell motility via its interaction with DOCK1 and DOCK4.,PTM:Phosphorylated on Tyr-221 upon cell adhesion. Results in the negative regulation of the association with SH2- and SH3- binding partners, possibly by the formation of an intramolecular interaction of phosphorylated Tyr-221 with the SH2 domain. This leads finally to the down-regulation of the Crk signaling pathway.,PTM:Phosphorylation of Crk-II (40 kDa) gives rise to a 42 kDa form.,similarity:Contains 1 SH2 domain.,similarity:Contains 1 SH3 domain.,similarity:Contains 2 SH3 domains.,subcellular location:Translocated to the plasma membrane upon cell adhesion.,subunit:Interacts with ABL1, C3G, SOS, MAP4K1, MAPK8 and DOCK3 via its first SH3 domain. Interacts with BCAR1, CBL, CBLB, PXN, IRS4 and GAB1 via its SH2 domain upon stimulus-induced tyrosine phosphorylation. Interacts with several tyrosine-phosphorylated growth factor receptors such as EGFR, PDGFR and INSR via its SH2 domain (By similarity). Interacts with DOCK1 and DOCK4. Interacts with SHB.,		

# Validation Data



Western Blot analysis using Crk II Monoclonal Antibody against HEK293 (1) and CRK-hIgGFc transfected HEK293 (2) cell lysate.



Immunohistochemistry analysis of paraffin-embedded intima cancer tissues with DAB staining using Crk II Monoclonal Antibody.



Immunofluorescence analysis of 3T3-L1 cells using Crk II Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye. Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



Flow cytometric analysis of Hela cells using Crk II Monoclonal Antibody (blue) and negative control (red).



## **Contact information**

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **Crk II Mouse mAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents