

Caspase-1 (PTR1317) Mouse mAb(HRP)

CatalogNo: YM2210

Key Features

Host Species

- Mouse

Reactivity

- Human, Mouse, Rat, Horse, Pig

Applications

- WB, IF, ELISA

MW

- 45kD (Observed)

Isotype

- IgG2b, Kappa

Conjugate

- HRP

Storage

Storage*

-15°C to -25°C/1 year(Do not lower than -25°C)

Formulation

PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Recommended Dilution Ratios

WB 1:500-2000

IF 1:100-500

ELISA 1:1000-5000

Basic Information

Clonality

Monoclonal

Clone Number

PTR1317

Immunogen Information

Immunogen

Synthesized peptide derived from human Caspase-1 AA range: 350-404

Specificity

This antibody detects endogenous levels of Caspase-1 protein.

Target Information

Gene name CASP1 IL1BC IL1BCE

Protein Name Caspase-1

Organism	Gene ID	UniProt ID
Human	834;	P29466;
Mouse	12362;	P29452;
Rat		P43527;

Cellular Localization Cytoplasm

Tissue specificity Expressed in larger amounts in spleen and lung. Detected in liver, heart, small intestine, colon, thymus, prostate, skeletal muscle, peripheral blood leukocytes, kidney and testis. No expression in the brain.

Function Thiol protease involved in a variety of inflammatory processes by proteolytically cleaving other proteins, such as the precursors of the inflammatory cytokines interleukin-1 beta (IL1B) and interleukin 18 (IL18) as well as the pyroptosis inducer Gasdermin-D (GSDMD), into active mature peptides . Plays a key role in cell immunity as an inflammatory response initiator: once activated through formation of an inflammasome complex, it initiates a pro-inflammatory response through the cleavage of the two inflammatory cytokines IL1B and IL18, releasing the mature cytokines which are involved in a variety of inflammatory processes . Cleaves a tetrapeptide after an Asp residue at position P1 . Also initiates pyroptosis, a programmed lytic cell death pathway, through cleavage of GSDMD . In contrast to cleavage of interleukins IL1B and IL18, recognition and cleavage of GSDMD is not strictly dependent on the consensus cleavage site but depends on an exosite interface on CASP1 that recognizes and binds the Gasdermin-D, C-terminal (GSDMD-CT) part . Upon inflammasome activation, during DNA virus infection but not RNA virus challenge, controls antiviral immunity through the cleavage of CGAS, rendering it inactive . In apoptotic cells, cleaves SPHK2 which is released from cells and remains enzymatically active extracellularly . ; [Isoform Delta]: Apoptosis inactive. ; [Isoform Epsilon]: Apoptosis inactive.

| Validation Data

| Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Caspase-1 (PTR1317) Mouse mAb(HRP)

