

RUNX2 (PT0524R) PT™ Rabbit mAb

CatalogNo: YM8347 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat, Human, Dog

Applications

- WB, IHC, IF, IP, ELISA

MW

- 57kD (Calculated)
- 57kD (Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

IHC 1:1000-1:5000

WB 1:2000-1:10000

IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

Basic Information

Clonality Monoclonal

Clone Number PT0524R

Immunogen Information

Specificity Endogenous

Target Information

Gene name RUNX2

Protein Name Runt-related transcription factor 2


Organism	Gene ID	UniProt ID
Human	860 ;	Q13950 ;
Mouse	12393 ;	Q08775 ;
Rat		Q9Z2J9 ;

Cellular Localization Nucleus


Tissue specificity Specifically expressed in osteoblasts.

Function Disease:Defects in RUNX2 are the cause of cleidocranial dysplasia (CCD) [MIM:119600]. CCD is an autosomal dominant skeletal disorder with high penetrance and variable expressivity. It is due to defective endochondral and intramembranous bone formation. Typical features include hypoplasia/aplasia of clavicles, patent fontanelles, wormian bones (additional cranial plates caused by abnormal ossification of the calvaria), supernumerary teeth, short stature, and other skeletal changes. In some cases defects in RUNX2 are exclusively associated with dental anomalies.,Domain:A proline/serine/threonine rich region at the C-terminus is necessary for transcriptional activation of target genes and contains the phosphorylation sites.,Function:Transcription factor involved in osteoblastic differentiation and skeletal morphogenesis. Essential for the maturation of osteoblasts and both intramembranous and endochondral ossification. CBF binds to the core site, 5'-PYGPYGGT-3', of a number of enhancers and promoters, including murine leukemia virus, polyomavirus enhancer, T-cell receptor enhancers, osteocalcin, osteopontin, bone sialoprotein, alpha 1(I) collagen, LCK, IL-3 and GM-CSF promoters (By similarity). Inhibits MYST4-dependent transcriptional activation.,PTM:Phosphorylated; probably by MAP kinases (MAPK) (By similarity). Isoform 3 is phosphorylated on Ser-340.,similarity:Contains 1 Runt domain.,subunit:Heterodimer of an alpha and a beta subunit. Interacts with HIVP3 (By similarity). The alpha subunit binds DNA as a monomer and through the Runt domain. DNA-binding is increased by heterodimerization. Interacts with G22P1 (Ku70) and XRCC5 (Ku80). Interacts with MYST3 and MYST4.,tissue specificity:Specifically expressed in osteoblasts.,

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-RUNX2 antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: MDA-MB-231 Lane 2: Saos-2 Lane 3: NIH-3T3 Lane 4: Rat womb Predicted band size: 57kDa Observed band size: 57kDa



Immunofluorescence analysis of Hela cell. 1, RUNX2 Antibody (red) was diluted at 1:200 (4° overnight). NSE Monoclonal Antibody (13E2) (green) was diluted at 1:200 (4° overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog: RS3611 was diluted at 1:1000 (room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 Catalog: RS3208 was diluted at 1:1000 (room temperature, 50min).



Human osteosarcoma was stained with anti-RUNX2 rabbit antibody



Mouse spleen was stained with anti-RUNX2 rabbit antibody

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
RUNX2 (PT0524R)
PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)