

p70 S6 Kinase (Phospho Thr389) (PT0721R) PT® Rabbit mAb

CatalogNo: YM8766 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IF, IP, ELISA

MW

- 59kD (Calculated)
- 70kD (Observed)

Isotype

- IgG, Kappa

Recommended Dilution Ratios

WB 1:2000-1:10000**IF 1:200-1:1000****ELISA 1:5000-1:20000****IP 1:50-1:200**

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Basic Information

Clonality Monoclonal**Clone Number** PT0721R


Immunogen Information

Specificity p70 S6 Kinase (Phospho Thr389) Monoclonal Antibody detects endogenous levels of Phospho Atm around the phosphorylation site of Thr389 protein. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites): GFTYV

Target Information

Gene name	RPS6KB1 STK14A P70S6K		
Protein Name	Ribosomal protein S6 kinase beta-1, p70 S6 Kinase (Thr389)		
	Organism	Gene ID	UniProt ID
	Human	6198 ;	P23443 ;
	Mouse	72508 ;	Q8BSK8 ;
	Rat	83840 ;	P67999 ;
Cellular Localization	Cell junction, synapse, synaptosome . Mitochondrion outer membrane. Mitochondrion. Colocalizes with URI1 at mitochondrion.; [Isoform Alpha I]: Nucleus. Cytoplasm.; [Isoform Alpha II]: Cytoplasm.		
Tissue specificity	Widely expressed.		
Function	Catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Activation by serine/threonine phosphorylation and protein kinase C, inactivated by type 2A phosphatase.,Function:Phosphorylates specifically ribosomal protein S6 in response to insulin or several classes of mitogens.,similarity:Belongs to the protein kinase superfamily.,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. S6 kinase subfamily.,similarity:Contains 1 AGC-kinase C-terminal domain.,similarity:Contains 1 protein kinase domain.,subunit:Interacts with PPP1R9A/neurabin-1.,tissue specificity:Widely expressed.,		

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-p70 S6 Kinase (Phospho Thr389) (PT0721R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: C6 Lane 2: C6 was starved with serum for 24 hours and treated with IGF-1(100ng/mL) for 20 minutes Lane 3: 3T3-L1 Lane 4: 3T3-L1 was starved overnight with serum and treated with IGF-1(100ng/mL) for 20 minutes Lane 5: HeLa Lane 6: HeLa was starved overnight with serum and treated with IGF-1(100ng/mL) for 20 minutes Predicted band size: 59kDa Observed band size: 70kDa

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: http://www.immunoway.com
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**p70 S6 Kinase
(Phospho Thr389)
(PT0721R) PT®
Rabbit mAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)