

# PAI-1 (PT1139R) PT® Rabbit mAb

CatalogNo: YM8871 Recombinant R

#### **Key Features**

**Host Species** 

Rabbit

Reactivity

Human, Mouse, Rat

**Applications** WB,IHC,IF,IP,ELISA

MW

 45kD (Calculated) 45kD (Observed)

Isotype

• IgG,Kappa

#### Recommended Dilution Ratios

IHC 1:200-1:1000 WB 1:2000-1:10000 IF 1:200-1:1000

ELISA 1:5000-1:20000

IP 1:50-1:200

### Storage

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

### **I** Basic Information

Clonality Monoclonal

**Clone Number** PT1139R

# Immunogen Information

**Specificity** Endogenous

## | Target Information

Gene name

SERPINE1

**Protein Name** 

Plasminogen activator inhibitor 1

Organism	Gene ID	UniProt ID
Human	<u>5054;</u>	<u>P05121;</u>
Mouse		<u>P22777;</u>
Rat	<u>24617;</u>	<u>P20961</u> ;


Cellular Localization Secreted.

**Tissue specificity** Expressed in endothelial cells (PubMed:2430793, PubMed:3097076). Found in plasma, platelets, and hepatoma and fibrosarcoma cells.


**Function** 

Disease: Defects in SERPINE1 are the cause of plasminogen activator inhibitor-1 deficiency (PAI-1 deficiency) [MIM:173360]. This deficiency is characterized by abnormal bleeding due to SERPINE1 defect in the plasma., Disease: High concentrations of SERPINE1 have been associated with thrombophilia [MIM:188050]; an autosomal dominant disorder in which affected individuals are prone to develop serious spontaneous thrombosis., Function: This inhibitor acts as 'bait' for tissue plasminogen activator, urokinase, and protein C. Its rapid interaction with TPA may function as a major control point in the regulation of fibrinolysis.,online information:Plasminogen activator inhibitor-1 entry,PTM:Inactivated by proteolytic attack of the urokinase-type (u-PA) and the tissue-type (TPA), cleaving the 369-Arg-I-Met-370 bond., similarity: Belongs to the serpin family, subunit: Interacts with VTN. Binds LRP1B; binding is followed by internalization and degradation., tissue specificity: Found in plasma and platelets and in endothelial, hepatoma and fibrosarcoma cells.,

### **Validation Data**



Human kidney was stained with anti-PAI-1 (PT1139R) Rabbit antibody



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-PAl-1 (PT1139R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: HUVEC-T1 Lane 2: SW480 Predicted band size: 45kDa Observed band size: 45kDa

#### **|** Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **PAI-1 (PT1139R)** 

PT® Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents