

Peroxiredoxin- 2 (PT1624R) PT™ Rabbit mAb

CatalogNo: YM9466 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, IF, ELISA

MW

- 22kD (Calculated)
- 22kD (Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

IHC 1:200-1:1000**WB 1:2000-1:10000****IF 1:200-1:1000****ELISA 1:5000-1:20000**

Basic Information

Clonality Monoclonal**Clone Number** PT1624R

Immunogen Information

Specificity Endogenous

Target Information

Gene name PRDX2

Protein Name Peroxiredoxin-2


Organism	Gene ID	UniProt ID
Human	7001 ;	P32119 ;
Mouse	21672 ;	Q61171 ;
Rat	29338 ;	P35704 ;

Cellular Localization Cytoplasm .


Tissue specificity Brain,Cajal-Retzius cell,Cerebellum,Colon carcinoma,Erythrocyte,Fetal brain cortex,Hypothal

Function Catalytic activity:2 R'-SH + ROOH = R'-S-S-R' + H(2)O + ROH.,Function:Involved in redox regulation of the cell. Reduces peroxides with reducing equivalents provided through the thioredoxin system. It is not able to receive electrons from glutaredoxin. May play an important role in eliminating peroxides generated during metabolism. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H(2)O(2).,miscellaneous:Inactivated upon oxidative stress by overoxidation of Cys-51 to Cys-SO(2)H and Cys-SO(3)H. Cys-SO(2)H is retroreduced to Cys-SOH after removal of H(2)O(2), while Cys-SO(3)H may be irreversibly oxidized.,miscellaneous:The active site is the redox-active Cys-51 oxidized to Cys-SOH. Cys-SOH rapidly reacts with Cys-172-SH of the other subunit to form an intermolecular disulfide with a concomitant homodimer formation. The enzyme may be subsequently regenerated by reduction of the disulfide by thioredoxin.,similarity:Belongs to the ahpC/TSA family.,similarity:Contains 1 thioredoxin domain.,subunit:Homodimer; disulfide-linked, upon oxidation. May be found as a toroid-shaped decamer composed of 5 dimers, depending on pH and calcium concentration. Interacts with TIPIN.,

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-Peroxiredoxin- 2 (PT1624R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: K562 Lane 2: C2C12 Lane 3: PC-12 Lane 4: Hela Predicted band size: 22kDa Observed band size: 22kDa



Human appendix was stained with anti-Peroxiredoxin- 2 (PT1624R) Rabbit antibody

| Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Peroxiredoxin- 2 (PT1624R) PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)