Applications

WB,ELISA



CHSTB Rabbit pAb

CatalogNo: YN0545

| Key Features

Host Species Reactivity

Rabbit
 Human, Mouse, Rat

MW Isotype
• 38kD (Observed) • IgG

Recommended Dilution Ratios

WB 1:500-2000 ELISA 1:5000-20000

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol,0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from part region of human protein

Specificity CHSTB Polyclonal Antibody detects endogenous levels of protein.

| Target Information

Gene name CHST11

Protein Name

Carbohydrate sulfotransferase 11 (Chondroitin 4-O-sulfotransferase 1) (Chondroitin 4sulfotransferase 1) (C4S-1) (C4ST-1) (C4ST1)

Organism	Gene ID	UniProt ID
Human	<u>50515;</u>	Q9NPF2;
Mouse		Q9JME2;
Rat		<u>P69478;</u>

Cellular Localization


Golgi apparatus membrane ; Single-pass type II membrane protein .

Tissue specificity Widely expressed. Highly expressed in spleen, thymus, bone marrow, peripheral blood leukocytes, lymph node, heart, brain, lung and placenta.

Function

Catalytic activity:3'-phosphoadenylyl sulfate + chondroitin = adenosine 3',5'-bisphosphate + chondroitin 4'-sulfate., Disease: A chromosomal aberration involving CHST11 is found in Bcell chronic lymphocytic leukemias. Translocation t(12;14)(g23;g32) with IgH., Function: Catalyzes the transfer of sulfate to position 4 of the N-acetylgalactosamine (GalNAc) residue of chondroitin. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage and is distributed on the surfaces of many cells and extracellular matrices. Can also sulfate Gal residues in desulfated dermatan sulfate. Preferentially sulfates in GlcA->GalNAc unit than in IdoA->GalNAc unit. Does not form 4, 6di-O-sulfated GalNAc when chondroitin sulfate C is used as an acceptor.,PTM:Nglycosylated; required for activity and stability., similarity: Belongs to the sulfotransferase 2 family., tissue specificity: Widely expressed. Highly expressed in spleen, thymus, bone marrow, peripheral blood leukocytes, lymph node, heart, brain, lung and placenta.,

I Validation Data



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night

I Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **CHSTB Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents