**Applications** 



# **GTR2 Rabbit pAb**

CatalogNo: YN1353

## **Key Features**

Host Species Reactivity

Rabbit
 Human,Rat,Mouse,
 WB,ELISA

MW Isotype
• 57kD (Observed) • IgG

### **Recommended Dilution Ratios**

WB 1:500-2000 ELISA 1:5000-20000

## Storage

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol,0.5% BSA and 0.02% sodium azide.

## **Basic Information**

**Clonality** Polyclonal

# Immunogen Information

**Immunogen** Synthesized peptide derived from human protein . at AA range: 220-300

**Specificity** GTR2 Polyclonal Antibody detects endogenous levels of protein.

# | Target Information

Gene name SLC2A2 GLUT2

#### **Protein Name**

Solute carrier family 2, facilitated glucose transporter member 2 (Glucose transporter type 2, liver) (GLUT-2)

Organism	Gene ID	UniProt ID
Human	<u>6514;</u>	<u>P11168;</u>
Mouse		<u>P14246;</u>
Rat		<u>P12336;</u>

### Cellular Localization


Cell membrane; Multi-pass membrane protein.

**Tissue specificity** Liver, insulin-producing beta cell, small intestine and kidney.

#### **Function**

Disease:Defects in SLC2A2 are the cause of Fanconi-Bickel syndrome (FBS) [MIM:227810]. FBS is a rare, well-defined clinical entity, inherited in an autosomal recessive mode and characterized by hepatorenal glycogen accumulation, proximal renal tubular dysfunction, and impaired utilization of glucose and galactose.,Function:Facilitative glucose transporter. This isoform likely mediates the bidirectional transfer of glucose across the plasma membrane of hepatocytes and is responsible for uptake of glucose by the beta cells; may comprise part of the glucose-sensing mechanism of the beta cell. May also participate with the Na(+)/glucose cotransporter in the transcellular transport of glucose in the small intestine and kidney.,online information:GLUT2 entry,PTM:N-glycosylated; required for stability and retention at the cell surface of pancreatic beta cells.,similarity:Belongs to the major facilitator superfamily. Sugar transporter (TC 2.A.1.1) family. Glucose transporter subfamily.,tissue specificity:Liver, insulin-producing beta cell, small intestine and kidney.,

## **I** Validation Data



Western Blot analysis of varius cell lysis. Primary Antibody was diluted at 1:1000. Secondary antibody(catalog#:RS23920 was diluted at 1:10000

## Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **GTR2 Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents