

PI 3 kinase p85 β (Phospho Tyr464) Rabbit pAb

CatalogNo: YP0225

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IHC, ELISA

MW

- 85kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

IHC 1:200-1:1000

WB 1:500-1:2000

ELISA 1:10000

Not yet tested in other applications.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized phospho-peptide around the phosphorylation site of human PI 3-kinase p85 β (phospho Tyr464)

Specificity

Phospho-PI 3-kinase p85 β (Y464) Polyclonal Antibody detects endogenous levels of PI 3-kinase p85 β protein only when phosphorylated at Y464. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):QLyEE

Target Information

Gene name PIK3R2

Protein Name Phosphatidylinositol 3-kinase regulatory subunit beta


Organism	Gene ID	UniProt ID
Human	5296 ;	O00459 ;
Mouse	18709 ;	O08908 ;
Rat		Q63788 ;

Cellular Localization nucleus,cytosol,phosphatidylinositol 3-kinase complex,


Tissue specificity Brain,Epithelium,Kidney,Placenta,

Function Function: Binds to activated (phosphorylated) protein-tyrosine kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane.,similarity: Belongs to the PI3K p85 subunit family.,similarity: Contains 1 Rho-GAP domain.,similarity: Contains 1 SH3 domain.,similarity: Contains 2 SH2 domains.,subunit: Heterodimer of a p110 (catalytic) and a p85 (regulatory) subunits.,


Validation Data



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PI 3-kinase p85 β (Phospho-Tyr464) Antibody



Western blot analysis of lysate from Jurkat cells, using phospho-PI 3-kinase p85 β (Phospho-Tyr464) antibody.



Immunohistochemical analysis of paraffin-embedded human Moderately differentiated hepatocellular carcinoma Antibody was diluted at 1:200(4 $^{\circ}$ overnight).

Contact information

Orders: order@immunoway.com
 Support: tech@immunoway.com
 Telephone: 877-594-3616 (Toll Free), 408-747-0185
 Website: <http://www.immunoway.com>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
PI 3 kinase p85 β (Phospho Tyr464) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)