

JNK2 Rabbit pAb

CatalogNo: YT2442

Key Features

Host Species

• Rabbit

Isotype

Reactivity

· Human, Mouse, Rat

• 48kD (Observed)

• IgG

WB,IHC,IF,ELISA

Applications

Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:10000 IF 1:50-200

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human MAPK9.

AA range:246-295

Specificity JNK2 Polyclonal Antibody detects endogenous levels of JNK2 protein.

Target Information

Gene name MAPK9

Protein Name Mitogen-activated protein kinase 9

Organism	Gene ID	UniProt ID
Human	<u>5601;</u>	<u>P45984;</u>
Mouse	<u>26420;</u> ;	Q9WTU6;
Rat	<u>50658;</u>	<u>P49186;</u>

Cellular Localization Cytoplasm . Nucleus . Colocalizes with POU5F1 in the nucleus. .

Tissue specificity Brain, Skin,

Function Catalytic activity:ATP + a protein = ADP + a

phosphoprotein.,cofactor:Magnesium.,Domain:The TXY motif contains the threonine and tyrosine residues whose phosphorylation activates the MAP kinases.,enzyme regulation:Activated by threonine and tyrosine phosphorylation by either of two dual specificity kinases, MAP2K4 and MAP2K7. Inhibited by dual specificity phosphatases, such as DUSP1.,Function:JNK2 isoforms display different binding patterns: alpha-1 and alpha-2 preferentially bind to c-Jun, whereas beta-1 and beta-2 bind to ATF2. However, there is no correlation between binding and phosphorylation, which is achieved at about the same efficiency by all isoforms. JUNB is not a substrate for JNK2 alpha-2, and JUND binds only weakly to it.,Function:Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as c-Jun and ATF2 and thus regulates AP-1 transcriptional activity. In T-cells, JNK1 and JNK2 are required for polarized differentiation of T-helper cells into Th1 cells.,PTM:Dually phosphorylated on Thr-183 and Tyr-185, which activates the enzyme. Autophosphorylated in vitro.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. MAP kinase subfamily, similarity:Contains 1, protein kinase

Ser/Thr protein kinase family. MAP kinase subfamily., similarity:Contains 1 protein kinase domain., subunit:Binds to at least four scaffolding proteins, MAPK8IP1/JIP-1, MAPK8IP2/JIP-2, MAPK8IP3/JIP-3/JSAP1 and SPAG9/MAPK8IP4/JIP-4. These proteins also bind other

components of the JNK signaling pathway. Interacts with NFATC4.,

Validation Data

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: JNK2 Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.	Antibody ELISA Kits Protein Reagents
	Immunoway - 3 /
	Immiinoway - 3 /