

IL-31R α Rabbit pAb

CatalogNo: YT5091

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, ELISA

MW

- 82kD (Observed)

Isotype

- IgG

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000**ELISA 1:40000****Not yet tested in other applications.**

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from the Internal region of human IL-31R α .**Specificity** IL-31R α Polyclonal Antibody detects endogenous levels of IL-31R α protein.

Target Information

Gene name IL31RA

Protein Name Interleukin-31 receptor subunit alpha


Organism	Gene ID	UniProt ID
Human	133396 ;	Q8NI17 ;
Mouse		Q8K5B1 ;

Cellular Localization Cell membrane ; Single-pass type I membrane protein . Cell junction, synapse, presynaptic cell membrane . Cell projection, axon .

Tissue specificity Expressed in CD14- and CD56-positive blood cells (PubMed:11877449). Expressed in macrophages (PubMed:16461143, PubMed:18439099). Expressed in keratinocytes (PubMed:21261663). Expressed in a subset of dorsal root ganglia neurons (at protein level) (PubMed:24373353). Expressed at low levels in testis, ovary, brain, prostate, placenta, thymus, bone marrow, trachea and skin (PubMed:15184896, PubMed:11877449, PubMed:14504285). Expressed in bronchial and alveolar epithelial cells and pulmonary fibroblasts (PubMed:18439099). Detected in all of the myelomonocytic lineage (PubMed:14504285). Isoform 6: Expressed at higher levels in lesional skin compared to healthy skin of atopic dermatitis patients (PubMed:24373353).

Function Function:Associates with OSMR to form the interleukin-31 receptor which activates STAT3 and to a lower extent STAT1 and STAT5. May function in skin immunity.,induction:Up-regulated in lesional keratinocytes of patients with atopic dermatitis. Up-regulated by IFNG and lipopolysaccharides.,PTM:N-glycosylated.,similarity:Belongs to the type I cytokine receptor family. Type 2 subfamily.,similarity:Contains 5 fibronectin type-III domains.,subunit:Heterodimer with OSMR. Interacts with JAK1 and STAT3.,tissue specificity:Expressed at low levels in testis, ovary, brain, prostate, placenta, thymus, bone marrow, trachea and skin. Detected in all of the myelomonocytic lineage. Expressed in CD14-and CD56-positive blood cells and by macrophages (at protein level),,

Validation Data



Western Blot analysis of extracts from Jurkat cells, using IL-31R Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000

| Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
IL-31R α Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)