

Goat Anti Human IgG gamma chain (FITC)

CatalogNo: RS2305

Key Features

Host Species
• Goat

Reactivity

Human

Applications
• IF,FC

Conjugate • FITC

Recommended Dilution Ratios

Optimal working dilutions should be determined experimentally by the investigator Suggested starting dilutions are:IF 1:250-1:2000 Flow Cyt 1:250-1:2000

Storage

Storage*	Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Immunogen Information

Target Information

Protein Name

Validation Data

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: Goat Anti Human IgG gamma chain (FITC)

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents