

Goat Anti Pig IgG(H+L) (FITC)

CatalogNo: RS2314

Key Features

Host Species

Goat

Reactivity

Pig

Applications

IF,FC

Conjugate

FITC

Recommended Dilution Ratios

Optimal working dilutions should be determined experimentally by the investigator Suggested starting dilutions are:IF 1:250-1:2000 Flow Cyt 1:250-1:2000

Storage

Storage*

Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Immunogen Information

| Target Information

Protein Name

| Validation Data

| Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

Goat Anti Pig
IgG(H+L) (FITC)

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents