

Goat Anti Rat IgG (DyLight 488)

CatalogNo: RS23240

| Key Features

Host Species

- Goat

Reactivity

- Rat

Applications

- IF,FC

Conjugate

- DyLight 488

| Recommended Dilution Ratios

Optimal working dilutions should be determined experimentally by the investigator
Suggested starting 1:50-1:1000 dilutions for most fluorescent applications.

| Storage

Storage*

Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing

Formulation

Liquid in PBS, pH 7.4, containing 0.02% Sodium Azide as preservative, 1% BSA as stabilizer and 50% Glycerol.

| Basic Information

| Immunogen Information

| Target Information

Protein Name

Validation Data

DyLight 350	353/432	Blue
DyLight 405	400/420	Blue
DyLight 488	493/518	Green
DyLight 549	562/576	Yellow
DyLight 594	593/618	Red/Orange
DyLight 649	652/672	Red
DyLight 680	692/712	Near IR
DyLight 800	777/794	Near IR

To use the DyLight Fluors with fluorescent imagers, use a spectral line of the blue laser diode for DyLight 405, a cyan (488 nm) laser for DyLight 488, a green (526 nm) laser for DyLight 550 and 594, and a red (633 nm) laser for DyLight 649. The DyLight 680 and 800 fluors are compatible with laser- and filter-based infrared imaging instruments that emit in the 700 nm and 800 nm

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Goat Anti Rat IgG (DyLight 488)

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)