

Goat Anti Mouse IgG (Dylight 800)

CatalogNo: RS23910

Key Features

Host Species

Goat

Reactivity Mouse

Applications IF,FC,WB

Coniugate

Dylight 800

Recommended Dilution Ratios

Suggested starting 1:50-1:1000 dilutions for most fluorescent applications while 1:5000-20000 dilutions for fluorescent assays with LI-1:COR® Odyssey equipment would be recommended.

Storage

Storage*

Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot

to avoid repeated freezing and thawing

Formulation

Liquid in PBS, pH 7.4, containing 0.02% Sodium Azide as preservative, 1% BSA as stablizer and 50% Glycerol.


I Basic Information

Immunogen Information


| Target Information

Protein Name

| Validation Data



Exosomes secreted from cardiomyocytes suppress the sensitivity of tumor ferroptosis in ischemic heart failure Signal Transduction and Targeted Therapy Baofeng Yang WB Mouse plasma LLC cell, K7M2 cell



To use the DyLight Fluors with fluorescent imagers, use a spectral line of the blue laser diode for DyLight 405, a cyan (488 nm) laser for DyLight 488, a green (526 nm) laser for DyLight 550 and 594, and a red (633 nm) laser for DyLight 649. The DyLight 680 and 800 fluors are compatible with laser- and filter-based infrared imaging instruments that emit in the 700 nm and 800 nm

| Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

Goat Anti Mouse IgG (Dylight 800)

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents