

Adiponectin Rabbit pAb

CatalogNo: YC0016

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA

MW

- 32kD (Observed)

Isotype

- IgG

Storage

Storage*

-15°C to -25°C/1 year (Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000**ELISA 1:5000****Not yet tested in other applications.**

Basic Information

Clonality

Polyclonal

Immunogen Information

Immunogen

Synthesized peptide derived from Adiponectin . at AA range: 10-90

Specificity

Adiponectin Polyclonal Antibody detects endogenous levels of Adiponectin protein.

Target Information

Gene name

ADIPOQ

Protein Name ADP/Acrp30 (Adiponectin)

Organism	Gene ID	UniProt ID
Human	9370 ;	Q15848 ;
Mouse	11450 ;	Q60994 ;

Cellular Localization Secreted .

Tissue specificity Synthesized exclusively by adipocytes and secreted into plasma.

Function Disease:Defects in ADIPOQ are the cause of adiponectin deficiency (ADPND) [MIM:612556]. ADPND results in very low concentrations of plasma adiponectin. ,Disease:Genetic variations in ADIPOQ are associated with non-insulin-dependent diabetes mellitus (NIDDM) [MIM:125853]; also known as diabetes mellitus type 2. NIDDM is characterized by an autosomal dominant mode of inheritance , onset during adulthood and insulin resistance. ,Domain:The C1q domain is commonly called the globular domain. ,Function:Important adipokine involved in the control of fat metabolism and insulin sensitivity , with direct anti-diabetic , anti-atherogenic and anti-inflammatory activities. Stimulates AMPK phosphorylation and activation in the liver and the skeletal muscle , enhancing glucose utilization and fatty-acid combustion. Antagonizes TNF-alpha by negatively regulating its expression in various tissues such as liver and macrophages , and also by counteracting its effects. Inhibits endothelial NF-kappa-B signaling through a cAMP-dependent pathway. May play a role in cell growth , angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities , depending on the type of complex , LMW , MMW or HMW. ,miscellaneous:HMW-complex blood contents are higher in females than in males , are increased in males by castration and decreased again upon subsequent testosterone treatment , which blocks HMW-complex secretion (By similarity) . In type 2 diabetic patients , both the ratios of HMW to total adiponectin and the degree of adiponectin glycosylation are significantly decreased as compared with healthy controls. ,miscellaneous:Variants Arg-84 and Ser-90 show impaired formation of HMW complexes whereas variants Cys-112 and Thr-164 show impaired secretion of adiponectin in any form. ,online information:Adiponectin entry ,pharmaceutical:Adiponectin might be used in the treatment of diabetes type 2 and insulin resistance. ,polymorphism:Genetic variations in ADIPOQ influence the variance in adiponectin serum levels and define the adiponectin serum levels quantitative trait locus 1 (ADIPQTL1) [MIM:612556]. ,PTM:HMW complexes are more extensively glycosylated than smaller oligomers. Hydroxylation and glycosylation of the lysine residues within the collagen-like domain of adiponectin seem to be critically involved in regulating the formation and/or secretion of HMW complexes and consequently contribute to the insulin-sensitizing activity of adiponectin in hepatocytes. ,PTM:Hydroxylated Lys-33 was not identified in PubMed:16497731 , probably due to poor representation of the N-terminal peptide in mass fingerprinting. ,PTM:Not N-glycosylated. ,PTM:O-linked glycans consist of Glc-Gal disaccharides bound to the oxygen atom of post-translationally added hydroxyl groups. ,similarity:Contains 1 C1q domain. ,similarity:Contains 1 collagen-like domain. ,subunit:Homomultimer. Forms trimers , hexamers and 12- to 18-mers. The trimers (low molecular weight complexes / LMW) are assembled via non-covalent interactions of the collagen-like domains in a triple helix and hydrophobic interactions within the globular C1q domain. Several trimers can associate to form disulfide-linked hexamers (middle molecular weight complexes / MMW) and larger complexes (higher molecular weight / HMW) . The HMW-complex assembly may rely additionally on lysine hydroxylation and glycosylation. LMW , MMW and HMW complexes bind to HBEGF , MMW and HMW complexes bind to PDGFB , and HMW complex binds to FGF2. ,tissue specificity:Synthesized exclusively by adipocytes and secreted into plasma. ,

| Validation Data

| Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
**Adiponectin Rabbit
pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)