

ACC1 α protein

CatalogNo: YD0001

| Key Features

Reactivity

- Human

Applications

- WB,SDS-PAGE

| Recommended Dilution Ratios

WB 1:500-2000

| Storage

Storage* -20°C/6 month,-80°C for long storage**Formulation** Liquid in PBS

| Basic Information

Source E.coli**Purification** E.coli**Purity** SDS-PAGE >90%

| Immunogen Information

Sequence Amino acid: 521-627, with his-MBP tag.

| Target Information

Protein Name ACC1 α protein**Organism**

Human

Gene ID[31](#);**UniProt ID**[O00763](#); [Q13085](#);


Cellular Localization

Mitochondrion .

Tissue specificity Widely expressed with highest levels in heart, skeletal muscle, liver, adipose tissue, mammary gland, adrenal gland and colon (PubMed:9099716). Isoform 3 is expressed in skeletal muscle, adipose tissue and liver (at protein level) (PubMed:19190759). Isoform 3 is detected at high levels in adipose tissue with lower levels in heart, liver, skeletal muscle and testis (PubMed:19190759).

Function fatty acid metabolic process, fatty acid biosynthetic process, lipid biosynthetic process, regulation of cellular ketone metabolic process, organic acid biosynthetic process, regulation of lipid metabolic process, regulation of fatty acid metabolic process, regulation of fatty acid oxidation, carboxylic acid biosynthetic process,

Validation Data



Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
ACC1 α protein

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)