

# Nrf2 protein

CatalogNo: YD0126

## **| Key Features**

Reactivity

**Applications**  Human WB,SDS-PAGE

### **Recommended Dilution Ratios**

WB 1:500-2000

## Storage

Storage\* -20°C/6 month,-80°C for long storage

**Formulation** Liquid in PBS

# **Basic Information**

**Source** E.coli

**Purification** E.coli

**Purity** SDS-PAGE >90%

# Immunogen Information

**Squence** Amino acid: 331-605, with his-MBP tag.

## | Target Information

NFE2L2 NRF2 **Gene name** 

#### **Protein Name**

Nrf2 protein

Organism	Gene ID	UniProt ID
Human	<u>4780</u> ;	<u>Q16236;</u>
Mouse	<u>18024;</u>	<u>Q60795</u> ;

### Cellular Localization


Cytoplasm, cytosol. Nucleus. Cytosolic under unstressed conditions: ubiquitinated and degraded by the BCR(KEAP1) E3 ubiquitin ligase complex (PubMed:15601839, PubMed:21196497). Translocates into the nucleus upon induction by electrophilic agents that inactivate the BCR(KEAP1) E3 ubiquitin ligase complex (PubMed:21196497). .

**Tissue specificity** Widely expressed. Highest expression in adult muscle, kidney, lung, liver and in fetal muscle.

#### **Function**

Domain: Acidic activation domain in the N-terminus, and DNA binding domain in the Cterminus., Function: Transcription activator that binds to antioxidant response (ARE) elements in the promoter regions of target genes. Important for the coordinated upregulation of genes in response to oxidative stress. May be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of hypersensitive site 2 of the beta-globin locus control region.,PTM:Phosphorylation of Ser-40 by PKC in response to oxidative stress dissociates NFE2L2 from its cytoplasmic inhibitor KEAP1, promoting its translocation into the nucleus., similarity: Belongs to the bZIP family., similarity: Belongs to the bZIP family. CNC subfamily., similarity: Contains 1 bZIP domain., subcellular location: Cytosolic under unstressed conditions, translocates into the nucleus upon induction by electrophilic agents., subunit: Heterodimer. May bind DNA with an unknown protein. Interacts with KEAP1. Interacts via its leucine-zipper domain with the coiled-coil domain of PMF1.,tissue specificity:Widely expressed. Highest expression in adult muscle, kidney, lung, liver and in fetal muscle.,

### Validation Data



### Contact information

Orders: order@immunoway.com tech@immunoway.com Support:

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the OR code to access additional product information:

Nrf2 protein

For Research Use Only. Not for Use in Diagnostic Procedures.	Antibody   ELISA Kits   Protein   Reagents
	Immunoway - 3 / 3