

# Progesterone receptor (PR) recombinant protein

CatalogNo: YD3022

## **Key Features**

Reactivity

Human

### Recommended Dilution Ratios

## Storage

**Storage\*** -15°C to -25°C/1 year(Avoid freeze / thaw cycles)

**Formulation** Phosphate-buffered solution

### **Basic Information**

Source Mammalian cells

Purification Mammalian cells

**Purity** >90% as determined by SDS-PAGE

Clone Number PR

# Immunogen Information

**Squence** Amino acid:165-305, with human FC tag.

### | Target Information

**Gene name** PGR;NR3C3

### **Protein Name**

Progesterone receptor (PR) (Nuclear receptor subfamily 3 group C member 3)

Organism	Gene ID	UniProt ID	
Human	<u>5241</u> ;	<u>P06401</u> ;	

### Cellular Localization

Nucleus. Cytoplasm. Note=Nucleoplasmic shuttling is both hormone- and cell cycledependent. On hormone stimulation, retained in the cytoplasm in the G(1) and G(2)/M phases.; [Isoform A]: Nucleus. Cytoplasm. Note=Mainly nuclear.; [Isoform 4]: Mitochondrion outer membrane.

**Tissue specificity** In reproductive tissues the expression of isoform A and isoform B varies as a consequence of developmental and hormonal status. Isoform A and isoform B are expressed in comparable levels in uterine glandular epithelium during the proliferative phase of the menstrual cycle. Expression of isoform B but not of isoform A persists in the glands during mid-secretory phase. In the stroma, isoform A is the predominant form throughout the cycle. Heterogeneous isoform expression between the glands of the endometrium basalis and functionalis is implying region-specific responses to hormonal stimuli.

### **Function**

The steroid hormones and their receptors are involved in the regulation of eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues. Depending on the isoform, progesterone receptor functions as a transcriptional activator or repressor.; [Isoform A]: Ligand-dependent transdominant repressor of steroid hormone receptor transcriptional activity including repression of its isoform B, MR and ER. Transrepressional activity may involve recruitment of corepressor NCOR2.; [Isoform B]: Transcriptional activator of several progesteron-dependent promoters in a variety of cell types. Involved in activation of SRC-dependent MAPK signaling on hormone stimulation.; [Isoform 4]: Increases mitochondrial membrane potential and cellular respiration upon stimulation by progesterone.

### **I** Validation Data

### Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

2200 Ringwood Ave San Jose, CA 95131 USA Address:



Please scan the QR code to access additional product information: **Progesterone** receptor (PR)

recombinant protein

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents