p57 (Acetyl Lys278) Rabbit pAb

CatalogNo: YK0044

Key Features

Host Species	Reactivity	Applications
• Rabbit	• Human,Mouse,Rat	• WB,ELISA

MW	Isotype
• 33kD (Observed)	• IgG

Storage

Storage*	-15°C to -25°C/1 year (Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Recommended Dilution Ratios

WB 1:500-1:2000
ELISA 1:20000
Not yet tested in other applications.

Basic Information

Clonality	Polyclonal

Immunogen Information

Immunogen	The antiserum was produced against synthesized Acetyl-peptide derived from human p57Kip2 around the Acetylation site of Lys278. AA range:241-290
Specificity	Acetyl-p57 (K278) Polyclonal Antibody detects endogenous levels of p57 protein only when acetylated at K278.The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):FAkRK

Target Information

Gene name CDKN1C

Protein Name Cyclin-dependent kinase inhibitor 1C


Organism	Gene ID	UniProt ID
Human	1028;	P49918;
Mouse	12577;	P49919;

Cellular Localization Nucleus .

Tissue specificity Expressed in the heart , brain , lung , skeletal muscle , kidney , pancreas and testis. Expressed in the eye. High levels are seen in the placenta while low levels are seen in the liver.

Function Disease:Defects in CDKN1C are a cause of Beckwith-Wiedemann syndrome (BWS) [MIM:130650]. BWS is a genetically heterogeneous disorder characterized by anterior abdominal wall defects including exomphalos (omphalocele) , pre- and postnatal overgrowth , and macroglossia. Additional less frequent complications include specific developmental defects and a predisposition to embryonal tumors. ,Disease:Defects in CDKN1C are involved in tumor formation. ,Function:Potent tight-binding inhibitor of several G1 cyclin/CDK complexes (cyclin E-CDK2 , cyclin D2-CDK4 , and cyclin A-CDK2) and , to lesser extent , of the mitotic cyclin B-CDC2. Negative regulator of cell proliferation. May play a role in maintenance of the non-proliferative state throughout life. ,similarity:Belongs to the CDI family. ,tissue specificity:Expressed in the heart , brain , lung , skeletal muscle , kidney , pancreas and testis. High levels are seen in the placenta while low levels are seen in the liver. ,

Validation Data



Western Blot analysis of 4T1 cells using Acetyl-p57 (K278) Polyclonal Antibody. Secondary antibody (catalog#:RS0002) was diluted at 1:20000

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: http://www.immunoway.com
Address: 2200 Ringwood Ave San Jose, CA 95131 USA

Please scan the QR code to access additional product information:
p57 (Acetyl Lys278) Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.