

#### www.immunoway.com

# **Bmx Mouse mAb**

CatalogNo: YM0076

# Key Features

Host Species

Mouse

Reactivity

Human

ApplicationsWB,ELISA

MW • 78kD (Calculated)

#### **Recommended Dilution Ratios**

WB 1:500-1:2000 ELISA 1:10000 Not yet tested in other applications.

### **Storage**

Storage\*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### **Basic Information**

Clonality Monoclonal

# Immunogen Information

Immunogen Purified recombinant fragment of human Bmx expressed in E. Coli.

**Specificity** Bmx Monoclonal Antibody detects endogenous levels of Bmx protein.

### **Target Information**

BMX

Gene name

#### **Protein Name** Cytoplasmic tyrosine-protein kinase BMX

	Organism	Gene ID	UniProt ID
	Human	<u>660;</u>	<u>P51813;</u>
	Mouse		<u>P97504;</u>
Cellular Localization	Cytoplasm . Localizes to the edges of spreading cells when complexed with BCAR1.		
Tissue specificity	Highly expressed in cells with great migratory potential, including endothelial cells and metastatic carcinoma cell lines.		
Function	Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,cofactor:Binds 1 zinc ion per subunit.,Domain:SH2 domain mediates interaction with RUFY1.,Function:Activity is required for interleukin 6 (IL-6) induced differentiation. May play a role in the growth and differentiation of hematopoietic cells. May be involved in signal transduction in endocardial and arterial endothelial cells.,induction:Activated by IL-6 through phosphatidylinositol 3-kinase (PI3-kinase) pathway. It is likely that activation occurs through binding of phosphoinositides to the PH domain.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. TEC subfamily.,similarity:Contains 1 Btk-type zinc finger.,similarity:Contains 1 PH domain.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 SH2 domain.,subunit:Interacts with RUFY1 and RUFY2.,tissue specificity:Preferentially expressed in epithelial and endothelial cells.,		

# Validation Data



Western Blot analysis using Bmx Monoclonal Antibody against BMX-hlgGFc transfected HEK293 cell lysate.

# **Contact information**

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **Bmx Mouse mAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents