

# c-Fgr Mouse mAb

CatalogNo: YM0151

## **Key Features**

**Host Species** 

Mouse

Reactivity

Human, Mouse

**Applications** 

WB,IHC,IF,ELISA

#### MW

59kD (Calculated)

# **Recommended Dilution Ratios**

WB 1:500-1:2000 IHC 1:200-1:1000 ELISA 1:10000 IF 1:50-200

### Storage

**Storage\*** -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### **Basic Information**

**Clonality** Monoclonal

#### Immunogen Information

**Immunogen** Purified recombinant fragment of human c-Fgr expressed in E. Coli.

**Specificity** c-Fgr Monoclonal Antibody detects endogenous levels of c-Fgr protein.

## **Target Information**

Gene name FGR SRC2

**Protein Name** Tyrosine-protein kinase Fgr

Organism	Gene ID	UniProt ID
Human	<u>2268</u> ;	<u>P09769;</u>
Mouse	<u>14191;</u>	<u>P14234;</u>

#### Cellular Localization


Cell membrane ; Lipid-anchor ; Cytoplasmic side . Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, ruffle membrane. Cytoplasm, cytosol. Cytoplasm, cytoskeleton. Mitochondrion inner membrane. Mitochondrion intermembrane space. Detected in mitochondrial intermembrane space and at inner membranes (By similarity). Colocalizes with actin fibers at membrane ruffles. Detected at plasma membrane lipid rafts.

**Tissue specificity** Detected in neutrophils, monocytes and natural killer cells (at protein level). Detected in monocytes and large lymphocytes.


**Function** 

Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. SRC subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 1 SH2 domain., similarity: Contains 1 SH3 domain., subunit: Binds PTPNS1.,

#### **| Validation Data**



Western Blot analysis using c-Fgr Monoclonal Antibody against HL60 (1) Raw264.7 (2) cell lysate.



Immunohistochemistry analysis of paraffin-embedded human Spleen tissues with AEC staining using c-Fgr Monoclonal Antibody.

#### | Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **c-Fgr Mouse mAb**