

IFN-γ Mouse mAb

CatalogNo: YM0354 Orthogonal Validated 💿

Key Features

Host Species

Reactivity
• Human

ApplicationsWB,ELISA

Mouse

MW

• 19kD (Calculated)

Recommended Dilution Ratios

WB 1:500-1:2000 ELISA 1:10000

Not yet tested in other applications.

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Monoclonal

Immunogen Information

Immunogen Recombinant human IFN-γ (BioSource company, Cat.No. PHC4033)

Specificity IFN-γ Monoclonal Antibody detects endogenous levels of IFN-γ protein.

| Target Information

Gene name IFNG

Protein Name

Interferon gamma

Organism	Gene ID	UniProt ID
Human	<u>3458;</u>	<u>P01579;</u>
Mouse		<u>P01580</u> ;

Cellular Localization


Secreted.

Tissue specificity Released primarily from activated T lymphocytes.

Function

Disease:In Caucasians, genetic variation in IFNG is associated with the risk of aplastic anemia (AA) [MIM:609135]. AA is a rare disease in which the reduction of the circulating blood cells results from damage to the stem cell pool in bone marrow. In most patients, the stem cell lesion is caused by an autoimmune attack. T-lymphocytes, activated by an endogenous or exogenous, and most often unknown antigenic stimulus, secrete cytokines, including IFN-gamma, which would in turn be able to suppress hematopoiesis., Function: Produced by lymphocytes activated by specific antigens or mitogens. IFN-gamma, in addition to having antiviral activity, has important immunoregulatory functions. It is a potent activator of macrophages, it has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor effects of the type I interferons., online information:Interferon gamma entry, pharmaceutical:Available under the name Actimmune (Genentech). Used for reducing the frequency and severity of serious infections associated with chronic granulomatous disease (CGD).,PTM:Proteolytic processing produces C-terminal heterogeneity, with proteins ending alternatively at Gly-150, Met-157 or Gly-161., similarity: Belongs to the type II (or gamma) interferon family,,subunit:Homodimer.,tissue specificity:Released primarily from activated T lymphocytes.,

Validation Data



Western Blot analysis using IFN-γ Monoclonal Antibody against IFN-γ recombinant protein.

| Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

IFN-y Mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents