

www.immunoway.com

MuSK Mouse mAb

CatalogNo: YM0457

Key Features

Host Species

Mouse

ReactivityHuman

Applications
• IHC,IF,ELISA

Recommended Dilution Ratios

IHC 1:200-1:1000 IF 1:200-1:1000 ELISA 1:10000 Not yet tested in other applications.

Storage

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Monoclonal

Immunogen Information

Immunogen	Purified recombinant extracellular fragment of human MuSK (aa24-209) fused with hlgGFc tag expressed in HEK293 cell line.
Specificity	MuSK Monoclonal Antibody detects endogenous levels of MuSK protein.

Target Information

Gene name MUSK

Ductoin Nouse	Mussla skale			Lines
Protein Name	Muscie, skele	etal receptor t	yrosine-protein	kinase

Organism	Gene ID	UniProt ID
Human	<u>4593;</u>	<u>015146;</u>
Mouse		<u>Q61006;</u>

- CellularCell junction, synapse, postsynaptic cell membrane ; Single-pass type I membrane protein .LocalizationColocalizes with acetylcholine receptors (AChR) to the postsynaptic cell membrane of the
neuromuscular junction. .
- FunctionCatalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine
phosphate.,Disease:Defects in MUSK is a cause of autosomal recessive congenital
myasthenic syndrome (CMS) [MIM:608931]. Congenital myasthenic syndromes are inherited
disorders of neuromuscular transmission that stem from mutations in presynaptic, synaptic,
or postsynaptic proteins. MUSK mutations lead to decreased agrin-dependent AChR
aggregation, a critical step in the formation of the neuromuscular
junction.,Function:Receptor tyrosine kinase that is a key mediator of agrin's action and is
involved in neuromuscular junction (NMJ) organization.,online information:MuSK
entry,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase
family.,similarity:Contains 1 FZ (frizzled) domain.,similarity:Contains 1 protein kinase
domain.,similarity:Contains 3 Ig-like C2-type (immunoglobulin-like)
domains.,subunit:Interacts with DOK7, which probably regulates its activity.,

Validation Data



Immunohistochemistry analysis of paraffin-embedded human lung cancer (A), muscles (B) and breast cancer (C) with DAB staining using MuSK Monoclonal Antibody.



Confocal immunofluorescence analysis of HEK293 cells trasfected with extracellular MUSK (aa24-209)-hlgGFc using MuSK Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye.

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **MuSK Mouse mAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents