

#### www.immunoway.com

# **PYK2 Mouse mAb**

CatalogNo: YM0543

## Key Features

Host Species

Mouse

Reactivity

Human

ApplicationsWB,IHC,IF,ELISA

MW • 116kD (Calculated)

### **Recommended Dilution Ratios**

WB 1:500-1:2000 IHC 1:200-1:1000 ELISA 1:10000 IF 1:50-200

#### **Storage**

Storage\*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

#### **Basic Information**

Clonality Monoclonal

#### Immunogen Information

Immunogen	Purified recombinant fragment of PYK2 (aa815-997) expressed in E. Coli.
Specificity	PYK2 Monoclonal Antibody detects endogenous levels of PYK2 protein.

#### **Target Information**

Gene name	PTK2B		
Protein Name	Protein-tyrosine kinase 2-beta <b>Organism</b>	Gene ID	UniProt ID
	Human	<u>2185;</u>	<u>Q14289;</u>
	Mouse		<u>Q9QVP9;</u>
Cellular Localization	Cytoplasm. Cytoplasm, perinuclear region. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell junction, focal adhesion. Cell projection, lamellipodium. Cytoplasm, cell cortex. Nucleus. Interaction with NPHP1 induces the membrane-association of the kinase. Colocalizes with integrins at the cell periphery.		
Tissue specificity	Most abundant in the brain, with high kidney (at protein level). Also expres		
Function	Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,Function:Involved in calcium induced regulation of ion channel and activation of the map kinase signaling pathway. May represent an important signaling intermediate between neuropeptide activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. Interacts with the SH2 domain of Grb2. May phosphorylate the voltage-gated potassium channel protein Kv1.2. Its activation is highly correlated with the stimulation of c-Jun N-terminal kinase activity. Involved in osmotic stress-dependent SNCA 'Tyr-125' phosphorylation.,PTM:Phosphorylated on tyrosine residues in response to various stimuli that elevate the intracellular calcium concentration, as well as by PKC activation. Recruitment by nephrocystin to cell matrix adhesions initiates Tyr-402 phosphorylation and kinase activation. Angiotensin II, thapsigargin and L-alpha-lysophosphatidic acid (LPA) also induce autophosphorylation and increase kinase activity.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family.,similarity:Contains 1 FERM domain.,similarity:Contains 1 protein kinase domain.,subcellular location:Interaction with nephrocystin induces the membrane- association of the kinase.,subunit:Interacts with Crk-associated substrate (Cas), PTPNS1 and SH2D3C (By similarity). Interacts with nephrocystin, ASAP2, OPHN1L, SKAP2 and TGFB111.,tissue specificity:Most abundant in the brain, with highest levels in amygdala and hippocampus. Low levels in kidney. Also expressed in spleen and lymphocytes.,		

# Validation Data



Western Blot analysis using PYK2 Monoclonal Antibody against Raji (1), PMA induced THP-1 (2), Jurkat (3) and Ramos (4) cell lysate.



Immunohistochemistry analysis of paraffin-embedded human Tonsil tissues with AEC staining using PYK2 Monoclonal Antibody.

## **Contact information**

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **PYK2 Mouse mAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents