

Rtn-3 Mouse mAb

CatalogNo: YM0568

| Key Features

Host Species

- Mouse

Reactivity

- Human

Applications

- IF, ELISA

| Recommended Dilution Ratios

IF 1:200-1:1000

ELISA 1:10000

Not yet tested in other applications

| Storage

Storage*

-15°C to -25°C/1 year(Do not lower than -25°C)

Formulation

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

| Basic Information

Clonality

Monoclonal

Clone Number

16C8

| Immunogen Information

Immunogen

Purified recombinant fragment of Rtn-3 expressed in E. Coli.

Specificity

Rtn-3 Monoclonal Antibody detects endogenous levels of Rtn-3 protein.


| Target Information

Gene name

RTN3

| | | | |
|------------------------------|--|------------------------|-------------------------|
| Protein Name | Reticulon-3 | | |
| | Organism | Gene ID | UniProt ID |
| | Human | 10313; | O95197; |
| | Mouse | | Q9ES97; |
| Cellular Localization | Endoplasmic reticulum membrane ; Multi-pass membrane protein . Golgi apparatus membrane ; Multi-pass membrane protein . | | |
| Tissue specificity | Isoform 3 is widely expressed, with highest levels in brain, where it is enriched in neuronal cell bodies from gray matter (at protein level). Three times more abundant in macula than in peripheral retina. Isoform 1 is expressed at high levels in brain and at low levels in skeletal muscle. Isoform 2 is only found in melanoma. | | |
| Function | <p>Function:May be involved in membrane trafficking in the early secretory pathway. Inhibits BACE1 activity and amyloid precursor protein processing. May induce caspase-8 cascade and apoptosis. May favor BCL2 translocation to the mitochondria upon endoplasmic reticulum stress. In case of enteroviruses infection, RTN3 may be involved in the viral replication or pathogenesis.,induction:By endoplasmic reticulum stress (at protein level).,miscellaneous:The sequence shown here is derived from an EMBL/GenBank/DDBJ third party annotation (TPA) entry.,similarity:Contains 1 reticulon domain.,subunit:Homodimer. Interacts with RTN4. Isoform 3 interacts with BACE1, BACE2, BCL2 and FADD. Interacts with Coxsackievirus A16, enterovirus 71 and poliovirus P2C proteins.,tissue specificity:Isoform 3 is widely expressed, with highest levels in brain, where it is enriched in neuronal cell bodies from gray matter (at protein level). Three times more abundant in macula than in peripheral retina. Isoform 1 is expressed at high levels in brain and at low levels in skeletal muscle. Isoform 2 is only found in melanoma.,</p> | | |

| Validation Data



Confocal immunofluorescence analysis of HeLa (A), A431 (B) and THP-1 (C) cells using Rtn-3 Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye.

| Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: http://www.immunoway.com
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
Rtn-3 Mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)