

## AIFM1 Mouse mAb

CatalogNo: YM1505

### Key Features

#### Host Species

- Mouse

#### Reactivity

- Human

#### Applications

- WB,IF

#### MW

- 67kD (Observed)

### Storage

**Storage\*** -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:1000**

**ICC 1:200**

**IF 1:50-200**

### Basic Information

**Clonality** Monoclonal

**Clone Number** 15E5

### Immunogen Information

**Immunogen** Purified recombinant human AIF protein fragments expressed in E.coli.

**Specificity** This antibody detects endogenous levels of AIF and does not cross-react with related proteins.

---

## | Target Information

**Gene name** AIFM1 ALF PDCD8

**Protein Name** Apoptosis-inducing factor 1 mitochondrial

Organism	Gene ID	UniProt ID
Human	<a href="#">9131</a> ;	<a href="#">O95831</a> ;
Mouse	<a href="#">26926</a> ;	<a href="#">Q9Z0X1</a> ;
Rat	<a href="#">83533</a> ;	<a href="#">Q9JM53</a> ;

**Cellular  
Localization**

Mitochondrion intermembrane space . Mitochondrion inner membrane. Cytoplasm . Nucleus . Cytoplasm, perinuclear region . Proteolytic cleavage during or just after translocation into the mitochondrial intermembrane space (IMS) results in the formation of an inner-membrane-anchored mature form (AIFmit). During apoptosis, further proteolytic processing leads to a mature form, which is confined to the mitochondrial IMS in a soluble form (AIFsol). AIFsol is released to the cytoplasm in response to specific death signals, and translocated to the nucleus, where it induces nuclear apoptosis (PubMed:15775970). Colocalizes with EIF3G in the nucleus and perinuclear region (PubMed:17094969). . ; [Isoform 3]: Mitochondrion intermembrane space . Mitochondrion inner membrane . Has a stronger membrane anchorage than isoform 1. . ; [Isoform 4]: Mitochondrion . Cytoplasm, cytosol . In pro-apoptotic conditions, is released from mitochondria to cytosol in a calpain/cathepsin-dependent manner. . ; [Isoform 5]: Cytoplasm .

**Tissue specificity** Expressed in all tested tissues (PubMed:16644725). Detected in muscle and skin fibroblasts (at protein level) (PubMed:23217327). Expressed in osteoblasts (at protein level) (PubMed:28842795). . ; [Isoform 3]: Brain specific. . ; [Isoform 4]: Expressed in all tested tissues except brain. . ; [Isoform 5]: Isoform 5 is frequently down-regulated in human cancers.

**Function**

Catalytic activity:2 glutathione + protein-disulfide = glutathione disulfide + protein-dithiol.,cofactor:FAD.,Function:Possesses significant protein thiol-disulfide oxidase activity.,Function:Probable oxidoreductase that acts as a caspase-independent mitochondrial effector of apoptotic cell death. Extramitochondrial AIF induces nuclear chromatin condensation and large scale DNA fragmentation (in vitro). Binds to DNA in a sequence-independent manner.,similarity:Belongs to the FAD-dependent oxidoreductase family.,similarity:Contains 1 thioredoxin domain.,subcellular location:Translocated to the nucleus upon induction of apoptosis.,subunit:Interacts with XIAP.,tissue specificity:Widely expressed.,

---

## | Validation Data

## | Contact information

Orders: order@immunoway.com  
Support: tech@immunoway.com  
Telephone: 877-594-3616 (Toll Free), 408-747-0185  
Website: http://www.immunoway.com  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**AIFM1 Mouse mAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)