

## GAPDH (2B8) Mouse mAb (FITC)

CatalogNo: YM2053

### Key Features

Host Species	Reactivity	Applications
• Mouse	• Human,Mouse,Rat,Mk,Dg,Ch,Hamster,Rabbit,Pig,sheep,Insect,Yeast	• WB,IF,IHC
Isotype • IgG1	Conjugate • FITC	

### Storage

**Storage\*** Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing. Store in dark.

**Formulation** Liquid in PBS, pH 7.4, containing 0.02% sodium azide as preservative and 50% Glycerol.

### Recommended Dilution Ratios

Optimal working dilutions should be determined experimentally by the investigator

Suggested starting dilutions are: IF 1:250-1:2000

Flow Cyt 1:250-1:2000

### Basic Information

**Clonality** Monoclonal

**Clone Number** 2B8

### Immunogen Information

**Specificity** GAPDH Monoclonal Antibody(2B8) FITC conjugated specially designed for your WB or IHC analysis.

### Target Information

<b>Gene name</b>	GAPDH				
<b>Protein Name</b>	Glyceraldehyde-3-phosphate dehydrogenase				
Organism	Gene ID	UniProt ID			
Human	<a href="#">2597</a> ;	<a href="#">P04406</a> ;			
Mouse	<a href="#">100042025</a> ;	<a href="#">P16858</a> ;			
Rat	<a href="#">24383</a> ;	<a href="#">P04797</a> ;			
<b>Cellular Localization</b>	Cytoplasm, cytosol . Nucleus . Cytoplasm, perinuclear region . Membrane . Cytoplasm, cytoskeleton . Translocates to the nucleus following S-nitrosylation and interaction with SIAH1, which contains a nuclear localization signal (By similarity). Postnuclear and Perinuclear regions (PubMed:12829261). .				
<b>Tissue specificity</b>	Astrocytoma, Brain, Cajal-Retzius cell, Colon adenocarcinoma, Epitheliu				
<b>Function</b>	Catalytic activity:D-glyceraldehyde 3-phosphate + phosphate + NAD(+) = 3-phospho-D-glyceroyl phosphate + NADH.,Function:Independent of its glycolytic activity it is also involved in membrane trafficking in the early secretory pathway.,online information:Glyceraldehyde 3-phosphate dehydrogenase entry,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 1.,pathway:Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 1/5.,PTM:Reversible S-nitrosylation of Cys-152 inhibits enzymatic activity and increases endogenous ADP-ribosylation, which inhibits the enzyme in a non-reversible manner. The latter modification is more likely to be a pathophysiological event associated with inhibition of gluconeogenesis.,sequence Caution:Differs quite extensively.,similarity:Belongs to the glyceraldehyde-3-phosphate dehydrogenase family.,subcellular location:Postnuclear and Perinuclear regions.,subunit:Homotetramer.,subunit:Homotetramer. Binds PRKCI.,				

## Validation Data

## Contact information

Orders: [order@immunoway.com](mailto:order@immunoway.com)  
 Support: [tech@immunoway.com](mailto:tech@immunoway.com)  
 Telephone: 877-594-3616 (Toll Free), 408-747-0185  
 Website: <http://www.immunoway.com>  
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**GAPDH (2B8) Mouse mAb (FITC)**