

β-Tubulin (5G3) Mouse mAb (AbFluor 594)

CatalogNo: YM2200

Key Features

Host Species Reactivity Applications
• Mouse • WB,IF,IHC

Human, Mouse, Rat, Mk, Dg, Ch, Hamster, Rabbit, sheep, Insect, Yeast

Isotype Conjugate
• IgG1 • AbFluor 594

Recommended Dilution Ratios

Optimal working dilutions should be determined experimentally by the investigator Suggested starting dilutions are as follows:IHC 1:200 IF 1:200.

Storage

Storage* Stable for one year at -15°C to -25°C from date of shipment. For maximum recovery of

product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot

to avoid repeated freezing and thawing. Store in dark.

Formulation Liquid in PBS, pH 7.4, containing 0.02% sodium azide as preservative and 50% Glycerol.

Basic Information

Clonality Monoclonal

Clone Number 5G3

Immunogen Information

Specificity β-Tubulin Monoclonal Antibody(5G3) AbFluor™ 594 Conjugated specially designed for

your Immunofluorescence analysis.

Target Information

Gene name

TUBB3

Protein Name

Tubulin beta-3 chain

Organism	Gene ID	UniProt ID
Human	<u>10381;</u>	<u>Q13509;</u>


Cellular Localization Cytoplasm, cytoskeleton. Cell projection, growth cone. Cell projection, lamellipodium. Cell projection, filopodium.

Tissue specificity Expression is primarily restricted to central and peripheral nervous system. Greatly increased expression in most cancerous tissues.

Function

Domain: The highly acidic C-terminal region may bind cations such as calcium., Function: Receptor for MSH (alpha, beta and gamma) and ACTH. The activity of this receptor is mediated by G proteins which activate adenylate cyclase..Function:Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alphachain.,polymorphism:Genetic variations in MC1R are associated with variation in skin/hair/eye pigmentation type 2 (SHEP2) [MIM:266300]. Hair, eye and skin pigmentation are among the most visible examples of human phenotypic variation, with a broad normal range that is subject to substantial geographic stratification. In the case of skin, individuals tend to have lighter pigmentation with increasing distance from the equator. By contrast, the majority of variation in human eye and hair color is found among individuals of European ancestry, with most other human populations fixed for brown eyes and black hair.,polymorphism:Variations in MC1R are linked to the degree of skin pigmentation (Types I-IV). Type I skin the most lightly pigmented and type IV the most dark pigmented. Partial loss-of-function mutations are associated with fair skin, poor tanning and increased skin cancer risk., similarity: Belongs to the G-protein coupled receptor 1 family., similarity: Belongs to the tubulin family., subunit: Dimer of alpha and beta chains., tissue specificity: Melanocytes and corticoadrenal tissue.,

Validation Data



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).

I Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: β-Tubulin (5G3) Mouse mAb (AbFluor 594)

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents