

Bcl-2 (PTR2303) Mouse mAb

CatalogNo: YM3041

KD/KO Validated **Orthogonal Validated** 

Key Features

Host Species

- Mouse

Reactivity

- Human, Mouse, Rat

Applications

- WB, IF, ELISA

MW

- 26kD (Calculated)
26kD (Observed)

Isotype

- IgG2b, Kappa

Recommended Dilution Ratios

WB 1:500-2000**IF 1:100-500****ELISA 1:1000-5000**

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Basic Information

Clonality Monoclonal**Clone Number** PTR2303

Immunogen Information

Immunogen Synthetic Peptide of human Bcl-2 AA range: 1-100**Specificity** This antibody detects endogenous levels of BCL-2 protein.

| Target Information

Gene name BCL2

Protein Name Apoptosis regulator Bcl-2


Organism	Gene ID	UniProt ID
Human	596;	P10415;
Mouse	12043;	P10417;
Rat	24224;	P49950;

Cellular Localization Membranous


Tissue specificity Expressed in a variety of tissues.

Function Disease:A chromosomal aberration involving BCL2 may be a cause of follicular lymphoma (FL) [MIM:151430]; also known as type II chronic lymphatic leukemia. Translocation t(14;18)(q32;q21) with immunoglobulin gene regions. BCL2 mutations found in non-Hodgkin lymphomas carrying the chromosomal translocation could be attributed to the Ig somatic hypermutation mechanism resulting in nucleotide transitions.,Domain:The BH4 motif is required for anti-apoptotic activity and for interaction with RAF-1.,Function:Suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells. Regulates cell death by controlling the mitochondrial membrane permeability. Appears to function in a feedback loop system with caspases. Inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1).,online information:Bcl-2 entry,PTM:Phosphorylation/dephosphorylation on Ser-70 regulates anti-apoptotic activity. Growth factor-stimulated phosphorylation on Ser-70 by PKC is required for the anti-apoptosis activity and occurs during the G2/M phase of the cell cycle. In the absence of growth factors, BCL2 appears to be phosphorylated by other protein kinases such as ERKs and stress-activated kinases. Dephosphorylated by protein phosphatase 2A (PP2A).,PTM:Proteolytically cleaved by caspases during apoptosis. The cleaved protein, lacking the BH4 motif, has pro-apoptotic activity, causes the release of cytochrome c into the cytosol promoting further caspase activity.,similarity:Belongs to the Bcl-2 family.,subunit:Forms homodimers, and heterodimers with BAX, BAD, BAK and Bcl-X(L). Heterodimerization with BAX requires intact BH1 and BH2 motifs, and is necessary for anti-apoptotic activity (By similarity). Also interacts with APAF1, RAF-1, TP53BP2, BBC3, BCL2L1, MRPL41 and BNIPL. Binding to FKBP8 seems to target BCL2 to the mitochondria and probably interferes with the binding of BCL2 to its targets.,tissue specificity:Expressed in a variety of tissues.,


| Validation Data




Various whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with anti-BCL-2(PTR2303)antibody. The HRP-conjugated Goat anti-Mouse IgG(H + L) antibody was used to detect the antibody. Lane 1: THP-1 Lane 2: MOLT-4 Lane 3: Raji Lane 4: HL-60 Lane 5: K562 Lane 6: Daudi Lane 7: Raw264.7 Lane 8: Mouse spleen Lane 9: Rat thymus



Western blot analysis of lysates from 1)HeLa cell , 2)HeLa cells knockdown by siRNA1 (F:GGAUGACUGAGUACCUGAATT,R:UUCAGGUACUCAGUCAUCCTT) siRNA2(F:GUGAUGAAGUACAUCUUAU,R:AUAAGGAUGUACUUAUCAC), (Green) primary antibody was diluted at 1:1000, 4° over night, Dylight 800 secondary antibody(Immunoway:RS23910)was diluted at 1:10000, 37° 1hour. (Red) GAPDH rabbit (Immunoway:YN5585) antibody was diluted at 1:5000 as loading control, 4° over night, Dylight 680 secondary antibody(Immunoway:RS23720)was diluted at 1:10000, 37° 1hour.



Zhang, J., Wong, C.C., Leung, K.T. et al. FGF18-FGFR2 signaling triggers the activation of c-Jun-YAP1 axis to promote carcinogenesis in a subgroup of gastric cancer patients and indicates translational potential. Oncogene 39, 6647-6663 (2020).



Wang, Peng, et al. "Resveratrol ameliorates autophagic flux to promote functional recovery in rats after spinal cord injury." Oncotarget 9.9 (2018): 8427.

Contact information

Orders: order@immunoway.com
 Support: tech@immunoway.com
 Telephone: 877-594-3616 (Toll Free), 408-747-0185
 Website: http://www.immunoway.com
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Bcl-2 (PTR2303)
Mouse mAb

