

CD79A (ABT-CD79A) Mouse mAb

CatalogNo: YM4790

Key Features

Host Species

- Mouse

Reactivity

- Human,Rat,

Applications

- IHC,WB,IF,ELISA

MW

- 25kD (Calculated)
- 39kD (Observed)

Isotype

- IgG2b,Kappa

Recommended Dilution Ratios

IHC 1:200-1000

WB 1:500-2000

IF 1:100-500

ELISA 1:1000-5000

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Basic Information

Clonality Monoclonal

Clone Number ABT-CD79a

Immunogen Information


Immunogen Synthesized peptide derived from human CD79A AA range: 100-226

Specificity The antibody can specifically recognize human CD79A protein.


| Target Information

Gene name	CD79A IGA MB1		
Protein Name	B-cell antigen receptor complex-associated protein alpha chain (Ig-alpha) (MB-1 membrane glycoprotein) (Membrane-bound immunoglobulin-associated protein) (Surface IgM-associated protein) (CD antigen CD79a)		
	Organism	Gene ID	UniProt ID
	Human	973 ;	P11912 ;
Cellular Localization	Cytoplasmic, Membranous		
Tissue specificity	B-cells.		
Function	<p>Disease:Defects in CD79A are a cause of non-Bruton type agammaglobulinemia [MIM:601495]. Agammaglobulinemia is an immunodeficiency disease which results in developmental defects in the maturation pathway of B-cells. Two different mutations, one at the splice donor site of intron 2 and the other at the splice acceptor site for exon 3, have been identified. Both mutations give rise to a truncated protein.,Function:Required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. Also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells. Stimulates SYK autophosphorylation and activation. Binds to BLNK, bringing BLNK into proximity with SYK and allowing SYK to phosphorylate BLNK. Also interacts with and increases activity of some Src-family tyrosine kinases. Represses BCR signaling during development of immature B cells.,online information:CD79A mutation db,PTM:Phosphorylated on tyrosine, serine and threonine residues upon B-cell activation. Phosphorylation of tyrosine residues by Src-family kinases is an early and essential feature of the BCR signaling cascade. The phosphorylated tyrosines serve as docking sites for SH2-domain containing kinases, leading to their activation which in turn leads to phosphorylation of downstream targets. Phosphorylation of serine and threonine residues may prevent subsequent tyrosine phosphorylation.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 ITAM domain.,subcellular location:Following antigen binding, the BCR has been shown to translocate from detergent-soluble regions of the cell membrane to lipid rafts although signal transduction through the complex can also occur outside lipid rafts.,subunit:Heterodimer of alpha and beta chains; disulfide-linked. Part of the B-cell antigen receptor complex where the alpha/beta chain heterodimer is non-covalently associated with an antigen-specific membrane-bound surface immunoglobulin of two heavy chains and two light chains. Interacts through its phosphorylated ITAM domain with the SH2 domains of SYK which stimulates SYK autophosphorylation and activation. Also interacts, when phosphorylated on Tyr-210, with the SH2 domain of BLNK/SLP65, bringing BLNK into proximity with SYK and allowing SYK to phosphorylate BLNK which is necessary for trafficking of the BCR to late endosomes. Interacts with Src-family tyrosine kinases including FYN and LYN, increasing their activity.,tissue specificity:B-cells.,</p>		


| Validation Data




Raji whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with anti-CD79a(ABT-CD79a) antibody. The HRP-conjugated Goat anti-Mouse IgG(H + L) antibody was used to detect the antibody. Lane 1: Raji




Human appendix tissue was stained with Anti-CD79a (ABT-CD79a) Antibody




Human lymphoma tissue was stained with Anti-CD79a (ABT-CD79a) Antibody




Human lymphoma tissue was stained with Anti-CD79a (ABT-CD79a) Antibody




Human spleen tissue was stained with Anti-CD79a (ABT-CD79a) Antibody



Human tonsil tissue was stained with Anti-CD79a (ABT-CD79a) Antibody



Rat spleen tissue was stained with Anti-CD79a (ABT-CD79a) Antibody



Rat spleen tissue was stained with Anti-CD79a (ABT-CD79a) Antibody

| Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
CD79A (ABT-CD79A)
Mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)