

# Nucleophosmin (ABT486) Mouse mAb

CatalogNo: YM4907

## Key Features

Host Species

Mouse

MW • 32kD (Calculated) 33kD (Observed) Reactivity

Human,Mouse,Rat,

Isotype

IgG2a,Kappa

Applications • IHC,WB,IF,ELISA

### Recommended Dilution Ratios

IHC 1:200-1000 WB 1:500-2000 IF 1:100-500 ELISA 1:1000-5000

### **Storage**

Storage\*-15°C to -25°C/1 year(Do not lower than -25°C)FormulationPBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

#### **Basic Information**

Clone Number ABT486

#### Immunogen Information

ImmunogenSynthesized peptide derived from human Nucleophosmin AA range: 100-200SpecificityThe antibody can specifically recognize human Nucleophosmin protein.

# Target Information

Gene name	NPM1 NPM		
Protein Name	Nucleophosmin <b>Organism</b>	Gene ID	UniProt ID
	Human	<u>4869;</u>	<u>P06748;</u>
Cellular Localization	Nuclear, Cytoplasmic		
Tissue specificity	Nuclear, Cytoplasmic		
Function	Disease:A chromosomal aberration involving NPM1 is a cause of myelodysplastic syndrome (MDS). Translocation t(3;5)(q25.1;q34) with MLF1.,Disease:A chromosomal aberration involving NPM1 is found in a form of acute promyelocytic leukemia. Translocation t(5;17)(q32;q11) with RARA.,Disease:A chromosomal aberration involving NPM1 is found in a form of non-Hodgkin lymphoma. Translocation t(2;5)(p23;q35) with ALK. The resulting chimeric NPM1-ALK protein homodimerize and the kinase becomes constitutively activated.,Disease:Defects in NPM1 are associated with acute myelogenous leukemia (AML). Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location.,Function:Involved in diverse cellular processes such as ribosome biogenesis, centrosome duplication, protein chaperoning, histone assembly, cell proliferation, and regulation of tumor suppressors TP53/p53 and ARF. Binds ribosome presumably to drive ribosome nuclear export. Associated with nucleolar ribonucleoprotein structures and bind single-stranded nucleic acids. Acts as a chaperonin for the core histones H3, H2B and H4.,PTM:Acetylated at C-terminal lysine residues, thereby increasing affinity to histones.,PTM:ADP-ribosylated.,PTM:Phosphorylated at Ser-4 by PLK1. Phosphorylated by CDK2 at Ser-125 and Thr-199. Phosphorylated at Ser-4 by PLK1. Phosphorylated by CDK2 at Ser-125 and Thr-199. Phosphorylated at Ser-70 by NEK2.,PTM:Sumoylated by ARF.,similarity:Belongs to the nucleoplasmin family.,subcellular location:Generally nucleolar, but is translocated to the nucleoplasm in case of serum starvation or treatment with anticancer drugs. Has been found in the cytoplasm in patients with primary acute myelogenous leukemia (AML), but not with secondary AML. Can shuttle between cytoplasm and nucleus.,subunit:Decamer formed by two pentameric rings associated in a head-to-head fashion. Disulfide-linked dimers under certain conditions. The SWAP complex consists of NPM1, NCL, PARP1 and SWAP70 (By similarity). Interacts with NSUN		

# Validation Data



A431 whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with anti-Nucleophosmin(ABT486)antibody. The HRP-conjugated Goat anti-Mouse IgG(H + L) antibody was used to detect the antibody. Lane 1: A431



Human colon carcinoma tissue was stained with Anti-Nucleophosmin (ABT486) Antibody



Human hepatocellular carcinoma tissue was stained with Anti-Nucleophosmin (ABT486) Antibody

### **Contact information**

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: Nucleophosmin (ABT486) Mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents