

LC3B (PT0235R) PT™ Rabbit mAb

CatalogNo: YM8147 Recombinant R

Key Features

Host Species

Rabbit

MW • 14kD,16kD (Calculated)

14kD,16kD (Observed)

Reactivity

Human, Mouse, Rat

Isotype

IgG,Kappa

Applications

WB,IHC,IF,ELISA

Recommended Dilution Ratios

IHC 1:200-1:1000 WB 1:1000-1:5000 IF 1:200-1:1000

ELISA 1:5000-1:20000

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

I Basic Information

Clonality Monoclonal

Clone Number PT0235R

Immunogen Information

Specificity Endogenous

| Target Information

Gene name MAP1LC3B

Protein Name MAP1LC3B

Organism Gene ID UniProt ID

3. ga		· · · · · · · · · · · · · · · · · · ·
Human	<u>81631</u> ;	Q9GZQ8;
Mouse		Q9CQV6;


Cellular Localization Cytoplasm

Tissue specificity Most abundant in heart, brain, skeletal muscle and testis. Little expression observed in liver.


Function

Caution:PubMed:12740394 has shown that the protein is cleaved at Lys-122 but PubMed:15355958 has shown that the cleavage site is at Gly-120 as in other mammalian orthologs.,Function:Probably involved in formation of autophagosomal vacuoles (autophagosomes).,PTM:The precursor molecule is cleaved by APG4B/ATG4B to form LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form LC3-II.,similarity:Belongs to the MAP1 LC3 family.,subcellular location:LC3-II binds to the autophagic membranes.,subunit:3 different light chains, LC1, LC2 and LC3, can associate with MAP1A and MAP1B proteins.,tissue specificity:Most abundant in heart, brain, skeletal muscle and testis. Little expression observed in liver.,


Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the primary antibody was used at 4°C, over night with a 1:5000 dilution . The Dylight 800-conjugated Goat anti-Rabbit antibody(Cat:RS23920) was used to detect the antibody. Lane1: HL60 - Human promyelocytic leukemia cell Lane2: HUVEC - Human umbilical vein endothelial cell Lane3: Human CAF - Human cancer-associated fibroblast Lane4: HCT116 - Human colorectal carcinoma Lane5: A549 - Human lung carcinoma Lane6: Jurkat - Human T lymphocyte leukemia Lane7: HT22 - Mouse hippocampal neuronal Lane8: Thp-1 - Human monocytic leukemia Lane9: HK-2 - Human proximal tubular epithelial Lane10: PC-12 - Rat adrenal pheochromocytoma Lane11: 3T3-L1 - Mouse embryonic fibroblast Lane12: PC-3 - Human prostate adenocarcinoma Lane13: U-14 - Mouse cervical carcinoma Lane14: 293F - HEK293 derivative, adapted for suspension culture Predicted band size: 14 16kDa Observed band size: 14 16kDa




Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-LC3B antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1:Jurkat Lane 2:Rat spleen Lane 3: Mouse brain Lane 3: U-87 MG Predicted band size: 14,16kDa Observed band size: 14,16kDa




Mouse brain was stained with anti-LC3B rabbit antibody




Rat brain was stained with anti-LC3B rabbit antibody




Human brain was stained with anti-LC3B rabbit antibody




Jia, Wei-Qiang, et al. "Verbascoside inhibits progression of glioblastoma cells by promoting Let-7g-5p and down-regulating HMGA2 via Wnt/beta-catenin signalling blockade." Journal of cellular and molecular medicine 24.5 (2020): 2901-2916.



Immunofluorescence analysis of Hela cell. 1,Cleaved-PARP-1 (D214) Antibody(red) was diluted at 1:200(4° overnight). LC3B Antibody(green) was diluted at 1:200(4° overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog:RS3611 was diluted at 1:1000(room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 Catalog:RS3208 was diluted at 1:1000(room temperature, 50min).




Immunofluorescence analysis of Mouse-lung tissue. 1,LC3B Antibody(red) was diluted at 1:200(4°C,overnight). 2, Cy3 labled Secondary antibody was diluted at 1:300(room temperature, 50min).3, Picture B: DAPI(blue) 10min. Picture A:Target. Picture B: DAPI. Picture C: merge of A+B



Feng, Qin. "Quantitative proteomic analysis reveals that Arctigenin alleviates concanavalin A-induced hepatitis through suppressing immune system and regulating autophagy." Frontiers in immunology 9 (2018): 1881.



rat brain was stained with anti-LC3B rabbit antibody



Immunofluorescence analysis of HEK293. Picture A: LC3B antibody (red). Picture B: DAPI (blue). Picture C: Merge of A+B

Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

LC3B (PT0235R) PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.	Antibody ELISA Kits Protein Reagents
	Immunoway - 5 / 5