

AMPK α1/2 (Phospho-Thr183/Thr172) (PT0920R) PT™ Rabbit mAb

CatalogNo: YM8689 Recombinant R

Key Features

Host Species

Rabbit

MW • 62kD,64kD (Calculated)

62kD (Observed)

Reactivity

· Human, Mouse, Rat

Isotype

IgG,Kappa

ApplicationsWB,IF,ELISA

Recommended Dilution Ratios

WB 1:2000-1:10000 IF 1:200-1:1000

ELISA 1:5000-1:20000

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Basic Information

Clonality Monoclonal

Clone Number PT0920R

Immunogen Information

Specificity

AMPK α 1/2 (Phospho-Thr183/Thr172) Antibody detects endogenous levels of Tau protein only when phosphorylated at Thr183/Thr172. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):LRtSC

| Target Information

Gene name

AAPK1/AAPK2

Protein Name

5'-AMP-activated protein kinase catalytic subunit alpha-1/2

Organism	Gene ID	UniProt ID
Human	<u>5562; 5563;</u>	Q13131; P54646;
Mouse	<u>105787; 108079;</u>	
Rat	65248; 78975;	<u>P54645; Q09137;</u>

Cellular Localization


Cytoplasm . Nucleus . In response to stress, recruited by p53/TP53 to specific promoters. .

Tissue specificity Brain,Intestine,Liver,Mammary gland,Platelet,Testis

Function

Catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:Binding of AMP results in allosteric activation, inducing phosphorylation on Thr-174 by STK11 in complex with STE20-related adapter-alpha (STRAD alpha) pseudo kinase and CAB39. Also activated by phosphorylation by CAMKK2 triggered by a rise in intracellular calcium ions, without detectable changes in the AMP/ATP ratio., Function: Responsible for the regulation of fatty acid synthesis by phosphorylation of acetyl-CoA carboxylase. It also regulates cholesterol synthesis via phosphorylation and inactivation of hormone-sensitive lipase and hydroxymethylglutaryl-CoA reductase. Appears to act as a metabolic stress-sensing protein kinase switching off biosynthetic pathways when cellular ATP levels are depleted and when 5'-AMP rises in response to fuel limitation and/or hypoxia. This is a catalytic subunit., sequence Caution: Translation N-terminally shortened., similarity: Belongs to the protein kinase superfamily., similarity: Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. SNF1 subfamily., similarity: Contains 1 protein kinase domain., subunit: Heterotrimer of an alpha catalytic subunit, a beta and a gamma non-catalytic subunits. Interacts with FNIP1 and FNIP2...

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-AMPK $\alpha1/2$ (Phospho-Thr183/Thr172) (PT0920R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: C6 Lane 2: C6 was treated oligomycin(0.5 μ M) for 30 minutes Predicted band size: 62,64kDa Observed band size: 62kDa

Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

AMPK α1/2 (Phospho-Thr183/Thr172) (PT0920R) PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents