

Src (Phospho Tyr529) (PT1540R) PT™ Rabbit mAb

CatalogNo: YM9382 **Recombinant** 

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, IF, IP, ELISA

MW

- 60kD (Calculated)
- 55kD (Observed)

Isotype

- IgG, Kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

WB 1:2000-1:10000**IF 1:200-1:1000****ELISA 1:5000-1:20000****IP 1:50-1:200**

Basic Information

Clonality Monoclonal**Clone Number** PT1540R

Immunogen Information


Immunogen The specific immunogen used to produce this antibody is proprietary information.

Specificity Src (Phospho Tyr529) (PT1540R) Antibody detects endogenous levels of c-Src protein only when phosphorylated at Y529. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):PQyQP

Target Information

Gene name	SRC SRC1		
Protein Name	Proto-oncogene tyrosine-protein kinase Src		
	Organism	Gene ID	UniProt ID
	Human	6714 ;	P12931 ;
	Mouse	20779 ;	P05480 ;
	Rat		Q9WUD9 ;
Cellular Localization	Cell membrane ; Lipid-anchor . Mitochondrion inner membrane . Nucleus . Cytoplasm , cytoskeleton . Cytoplasm , perinuclear region . Cell junction , focal adhesion . Localizes to focal adhesion sites following integrin engagement (PubMed:22801373) . Localization to focal adhesion sites requires myristoylation and the SH3 domain (PubMed:7525268) . Colocalizes with PDLIM4 at the perinuclear region , but not at focal adhesions (PubMed:19307596) . .		
Tissue specificity	Expressed ubiquitously. Platelets , neurons and osteoclasts express 5-fold to 200-fold higher levels than most other tissues.; [Isoform 1]: Expressed in spleen and liver. ; [Isoform 2]: Expressed in brain. ; [Isoform 3]: Expressed in brain.		
Function	Catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. ,PTM:Phosphorylated on Tyr-530 by c-Src kinase (CSK) . The phosphorylated form is termed pp60c-src. The phosphorylated tail interacts with the SH2 domain thereby repressing kinase activity. ,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. SRC subfamily. ,similarity:Contains 1 protein kinase domain. ,similarity:Contains 1 SH2 domain. ,similarity:Contains 1 SH3 domain. ,subunit:Interacts with DDEF1/ASAP1; via the SH3 domain. Interacts with CCPG1 (By similarity) . Interacts with CDCP1 , PELP1 , TGFBI1 and TOM1L2. Interacts with the cytoplasmic domain of MUC1 , phosphorylates it and increases binding of MUC1 with beta-catenin. Interacts with RALGPS1; via the SH3 domain. Interacts with HEV ORF3 protein; via the SH3 domain. ,		

Validation Data



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-Src (Phospho Tyr529) (PT1540R) antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: A431 Predicted band size: 60kDa Observed band size: 55kDa

Contact information

Orders: order@immunoway.com
 Support: tech@immunoway.com
 Telephone: 877-594-3616 (Toll Free), 408-747-0185
 Website: <http://www.immunoway.com>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
Src (Phospho Tyr529) (PT1540R) PT™ Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)