

IPO5 Rabbit pAb

CatalogNo: YN0803

| Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, ELISA

MW

- 120kD (Observed)

Isotype

- IgG

| Recommended Dilution Ratios

WB 1:500-2000

ELISA 1:5000-20000

| Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

| Basic Information

Clonality Polyclonal

| Immunogen Information

Immunogen Synthesized peptide derived from part region of human protein

Specificity IPO5 Polyclonal Antibody detects endogenous levels of protein.

| Target Information

Gene name IPO5 KPNB3 RANBP5

Protein Name	Importin-5 (Imp5) (Importin subunit beta-3) (Karyopherin beta-3) (Ran-binding protein 5) (RanBP5)		
	Organism	Gene ID	UniProt ID
	Human	3843 ;	O00410 ;
	Mouse		Q8BKC5 ;
Cellular Localization	Cytoplasm. Nucleus. Nucleus, nucleolus. Nucleus; nuclear rim. Found particularly in the nuclear rim and nucleolus.		
Tissue specificity	Bone marrow,Epithelium,Lung,Placenta,T-cell,Thalamus,		
Function	<p>Function:Functions in nuclear protein import as nuclear transport receptor. Serves as receptor for nuclear localization signals (NLS) in cargo substrates. Is thought to mediate docking of the importin/substrate complex to the nuclear pore complex (NPC) through binding to nucleoporin and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to the importin, the importin/substrate complex dissociates and importin is re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus (By similarity). Mediates the nuclear import of ribosomal proteins RPL23A, RPS7 and RPL5. Binds to a beta-like import receptor binding (BIB) domain of RPL23A. In vitro, mediates nuclear import of H2A, H2B, H3 and H4 histones. In case of HIV-1 infection, binds and mediates the nuclear import of HIV-1 Rev.,similarity:Belongs to the importin beta family.,similarity:Contains 1 importin N-terminal domain.,similarity:Contains 6 HEAT repeats.,subcellular location:Nucleus; nuclear rim. Found particularly in the nuclear rim and nucleolus.,subunit:Binds RPL23A, RPS7 and RPL5. Interacts with H2A, H2B, H3 and H4 histones (By similarity). Binds to HIV-1 Rev.,</p>		

| Validation Data

| Contact information

Orders:

order@immunoway.com

Support:

tech@immunoway.com

Telephone:

877-594-3616 (Toll Free), 408-747-0185

Website:

http://www.immunoway.com

Address:

2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

IPO5 Rabbit pAb