

## IPO9 Rabbit pAb

CatalogNo: YN0805

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse

#### Applications

- WB, ELISA

#### MW

- 114kD (Observed)

#### Isotype

- IgG

### Recommended Dilution Ratios

WB 1:500-2000

ELISA 1:5000-20000

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** Synthesized peptide derived from part region of human protein

**Specificity** IPO9 Polyclonal Antibody detects endogenous levels of protein.

### Target Information

**Gene name** IPO9 IMP9 KIAA1192 RANBP9 HSPC273

<b>Protein Name</b>	Importin-9 (Imp9) (Ran-binding protein 9) (RanBP9)		
	<b>Organism</b>	<b>Gene ID</b>	<b>UniProt ID</b>
	Human	<a href="#">55705;</a>	<a href="#">Q96P70;</a>
	Mouse		<a href="#">Q91YE6;</a>
<b>Cellular Localization</b>	Cytoplasm . Nucleus .		
<b>Tissue specificity</b>	Amygdala,B-cell lymphoma,Brain,Placenta,Umbilical cord blood,		
<b>Function</b>	<p>Function:Functions in nuclear protein import as nuclear transport receptor. Serves as receptor for nuclear localization signals (NLS) in cargo substrates. Is thought to mediate docking of the importin/substrate complex to the nuclear pore complex (NPC) through binding to nucleoporin and the complex is subsequently translocated through the pore by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to the importin, the importin/substrate complex dissociates and importin is re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus (By similarity). Mediates the nuclear import of H2B histone (By similarity), RPS7 and RPL18A. Prevents the cytoplasmic aggregation of RPS7 and RPL18A by shielding exposed basic domains. May also import H2A, H3, H4 histones (By similarity), RPL4 and RPL6.,similarity:Belongs to the importin beta family.,similarity:Contains 1 importin N-terminal domain.,subunit:Binds with high affinity to RPS7 and RPL18A. The binding is coupled to RanGTP cycles. May bind H2A, H3, H4 histones (By similarity), RPL4 and RPL6 with low affinity. Interacts with PPP2R1A and PPP2R1B.,</p>		

| Validation Data

| Contact information

Orders:
Support:
Telephone:
Website:
Address:

order@immunoway.com
tech@immunoway.com
877-594-3616 (Toll Free), 408-747-0185
http://www.immunoway.com
2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
**IPO9 Rabbit pAb**