

MIA3 Rabbit pAb

CatalogNo: YN0850

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse

Applications

- WB, ELISA

MW

- 209kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-2000

ELISA 1:5000-20000

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human protein . at AA range: 1110-1190

Specificity MIA3 Polyclonal Antibody detects endogenous levels of protein.

Target Information

Gene name MIA3 KIAA0268 TANGO UNQ6077/PRO20088

Protein Name Melanoma inhibitory activity protein 3 (C219-reactive peptide) (D320) (Transport and Golgi organization protein 1) (TANGO1)

Organism	Gene ID	UniProt ID
Human	375056 ;	Q5JRA6 ;
Mouse		Q8BI84 ;

Cellular Localization Endoplasmic reticulum membrane ; Single-pass membrane protein . Localizes at endoplasmic reticulum exit sites (ERES), also known as transitional endoplasmic reticulum (tER) (PubMed:32101163). SEC16A is required for its proper localization to ERES. After loading of COL7A1 into transport carriers, it is not incorporated into COPII carriers and remains in the endoplasmic reticulum membrane. .

Tissue specificity Broadly expressed, except in bone marrow and peripheral blood mononuclear cells. Down-regulated in melanoma tissue.

Function Domain:Although 2 transmembrane domains are predicted, PubMed:19269366 showed that it only contains one transmembrane domain. The other predicted transmembrane region is probably a hairpin-type region embedded into the membrane, which does not cross the membrane. It is unclear which of the 2 predicted transmembrane regions is the transmembrane or the hairpin-type region.,Domain:The proline-rich region (PRD) mediates the interaction with COPII coat subunits Sec23/24.,Function:Required for collagen VII (COL7A1) secretion by loading COL7A1 into transport carriers. May participate in cargo loading of COL7A1 at endoplasmic reticulum exit sites by binding to COPII coat subunits Sec23/24 and guiding SH3-bound COL7A1 into a growing carrier. Does not play a role in global protein secretion and is apparently specific to COL7A1 cargo loading. However, it may participate in secretion of other proteins in cells that do not secrete COL7A1.,similarity:Belongs to the MIA/OTOR family. Tango1 subfamily.,similarity:Contains 1 SH3 domain.,subcellular location:Localizes at endoplasmic reticulum exit sites. After loading of COL7A1 into transport carriers, it is not incorporated into COPII carriers and remains in the endoplasmic reticulum membrane.,subunit:Interacts (via SH3 domain) with COL7A1. Associates with the COPII coat subunits Sec23/Sec24.,tissue specificity:Broadly expressed, except in bone marrow and peripheral blood mononuclear cells. Down-regulated in melanoma tissue.,

| Validation Data

| Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
MIA3 Rabbit pAb

