

PR40A Rabbit pAb

CatalogNo: YN1097

Key Features

Host Species
• Rabbit
MW

Reactivity
• Human,Mouse
Isotype

• IgG

ApplicationsWB,ELISA

Recommended Dilution Ratios

WB 1:500-2000 ELISA 1:5000-20000

105kD (Observed)

Storage

| Storage* | -15°C to -25°C/1 year(Do not lower than -25°C) |
|-------------|--|
| Formulation | Liquid in PBS containing 50% glycerol,0.5% BSA and 0.02% sodium azide. |

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human protein . at AA range: 450-530

Specificity PR40A Polyclonal Antibody detects endogenous levels of protein.

Target Information

Gene name PRPF40A FBP11 FLAF1 FNBP3 HIP10 HYPA HSPC225

Protein Name Pre-mRNA-processing factor 40 homolog A (Fas ligand-associated factor 1) (Formin-binding protein 11) (Formin-binding protein 3) (Huntingtin yeast partner A) (Huntingtin-interacting protein 10) (HIP-10) (Huntingtin-interacting protein A) (Renal carcinoma antigen NY-REN-6)

| Organism | Gene ID | UniProt ID | |
|----------|---------------|----------------|--|
| Human | <u>55660;</u> | <u>075400;</u> | |
| Mouse | | <u>Q9R1C7;</u> | |

Cellular Nucleus speckle . Nucleus matrix . Colocalizes with AKAP8L in the nuclear matrix . Localization

Tissue specificity Expressed in the brain cortex (at protein level). Widely expressed.

Function Domain: The WW domains are essential for localization to nuclear speckles., Function: Binds to WASL/N-WASP and suppresses its translocation from the nucleus to the cytoplasm, thereby inhibiting its cytoplasmic function (By similarity). May be involved in pre-mRNA splicing., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence Caution: Contaminating sequence. Potential poly-A sequence starting in position 409., sequence caution: Contaminating sequence starting in position 409., sequence starting is position 400., sequence starting is position 400. Sequence starting is

Validation Data

Contact information

| Orders: | order@immunoway.com |
|------------|--|
| Support: | tech@immunoway.com |
| Telephone: | 877-594-3616 (Toll Free), 408-747-0185 |
| Website: | http://www.immunoway.com |
| Address: | 2200 Ringwood Ave San Jose, CA 95131 USA |
| | |



Please scan the QR code to access additional product information: **PR40A Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents