

RPGF2 Rabbit pAb

CatalogNo: YN1147

Key Features

Host Species

Rabbit

Reactivity

Human,Rat,Mouse

ApplicationsWB,ELISA

MW • 164kD (Observed) Isotype • IgG

Recommended Dilution Ratios

WB 1:500-2000 ELISA 1:5000-20000

Storage

| Storage* | -15°C to -25°C/1 year(Do not lower than -25°C) |
|-------------|--|
| Formulation | Liquid in PBS containing 50% glycerol,0.5% BSA and 0.02% sodium azide. |

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human protein . at AA range: 190-270

Specificity RPGF2 Polyclonal Antibody detects endogenous levels of protein.

Target Information

Gene name RAPGEF2 KIAA0313 NRAPGEP PDZGEF1

Protein Name

Rap guanine nucleotide exchange factor 2 (Neural RAP guanine nucleotide exchange protein) (nRap GEP) (PDZ domain-containing guanine nucleotide exchange factor 1) (PDZ-GEF1) (RA-GEF)

| Organism | Gene ID | UniProt ID |
|----------|--------------|----------------|
| Human | <u>9693;</u> | <u>Q9Y4G8;</u> |
| Mouse | | <u>Q8CHG7;</u> |
| Rat | | <u>F1M386;</u> |

- Cellular Localization Cytoplasm. Cytoplasm, perinuclear region. Cell membrane. Late endosome. Cell junction . Associated with the synaptic plasma membrane. Colocalizes with ADRB1 at the plasma membrane. Synaptosome. Enriched in synaptic plasma membrane and neuronal cell body. Colocalized with CTNNB1 at cell-cell contacts (By similarity). Localized diffusely in the cytoplasm before neuronal growth factor (NGF) stimulation. Recruited to late endosomes after NGF stimulation. Colocalized with the high affinity nerve growth factor receptor NTRK1 at late endosomes. Translocated to the perinuclear region in a RAP1A-dependent manner. Translocated to the cell membrane. .
- **Tissue specificity** Expressed in primary neuronal and endocrine cells (at protein level). Highest expression levels in brain. Lower expression levels in heart, kidney, lung, placenta and blood leukocytes.
- Function Function:Guanine nucleotide exchange factor (GEF) for Rap1A, Rap1B and Rap2B GTPases. Does not interact with cAMP or cGMP.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 cyclic nucleotide-binding domain.,similarity:Contains 1 Nterminal Ras-GEF domain.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 Ras-associating domain.,similarity:Contains 1 Ras-GEF domain.,subcellular location:Associated with the synaptic plasma membrane.,subunit:Interacts with MAGI2.,tissue specificity:Highest expression levels in brain. Lower expression levels in heart, kidney, lung, and placenta.,

Validation Data

Contact information

| Orders: | order@immunoway.com |
|------------|--|
| Support: | tech@immunoway.com |
| Telephone: | 877-594-3616 (Toll Free), 408-747-0185 |
| Website: | http://www.immunoway.com |
| Address: | 2200 Ringwood Ave San Jose, CA 95131 USA |



Please scan the QR code to access additional product information: **RPGF2 Rabbit pAb**