

# **RASL2 Rabbit pAb**

CatalogNo: YN1153

# **Key Features**

Host Species Reactivity Applications
• Rabbit • Human, Mouse • WB, ELISA

MW Isotype • 88kD (Observed) • IgG

### **Recommended Dilution Ratios**

WB 1:500-2000 ELISA 1:5000-20000

## Storage

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol,0.5% BSA and 0.02% sodium azide.

### **Basic Information**

**Clonality** Polyclonal

# Immunogen Information

**Immunogen** Synthesized peptide derived from human protein . at AA range: 100-180

**Specificity** RASL2 Polyclonal Antibody detects endogenous levels of protein.

## | Target Information

**Gene name** RASA4 CAPRI GAPL KIAA0538

#### **Protein Name**

Ras GTPase-activating protein 4 (Calcium-promoted Ras inactivator) (Ras p21 protein activator 4) (RasGAP-activating-like protein 2)

Organism	Gene ID	UniProt ID
Human	<u>10156;</u>	<u>043374;</u>
Mouse		Q6PFQ7;

### Cellular Localization

Cytoplasm, cytosol . Cell membrane ; Peripheral membrane protein . Localized to the cytosol as a result of its lack of phosphoinositide binding activity. Upon agonist-stimulated calcium mobilization, utilizes the C2A and C2B domains to associate with the plasma membrane.

### **Tissue specificity** Widely expressed.

#### **Function**

Domain:The PH domain does not bind phosphatidylinositol 4,5-bisphosphate or phosphatidylinositol 3,4,5-triphosphate. This lack of binding activity is due to Leu-592, compared to Arg found in other family members.,Function:Ca(2+)-dependent Ras GTPase-activating protein, that switches off the Ras-MAPK pathway following a stimulus that elevates intracellular calcium. Functions as an adaptor for Cdc42 and Rac1 during FcR-mediated phagocytosis.,similarity:Contains 1 Btk-type zinc finger.,similarity:Contains 1 PH domain.,similarity:Contains 1 Ras-GAP domain.,similarity:Contains 2 C2 domains.,subcellular location:Localized to the cytosol as a result of its lack of phosphoinositide binding activity. Upon agonist-stimulated calcium mobilization, utilizes the C2A and C2B domains to associate with the plasma membrane.,tissue specificity:Widely expressed.,

### **Validation Data**

# | Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **RASL2 Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents