

## SH3K1 Rabbit pAb

CatalogNo: YN1307

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human,Rat,Mouse

#### Applications

- WB,ELISA

#### MW

- 73kD (Observed)

#### Isotype

- IgG

### Recommended Dilution Ratios

**WB 1:500-2000**

**ELISA 1:5000-20000**

### Storage

**Storage\*** -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol,0.5% BSA and 0.02% sodium azide.

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** Synthesized peptide derived from part region of human protein

**Specificity** SH3K1 Polyclonal Antibody detects endogenous levels of protein.

### Target Information

**Gene name** SH3KBP1 CIN85

<b>Protein Name</b>	SH3 domain-containing kinase-binding protein 1 (CD2-binding protein 3) (CD2BP3) (Cbl-interacting protein of 85 kDa) (Human Src family kinase-binding protein 1) (HSB-1)		
	<b>Organism</b>	<b>Gene ID</b>	<b>UniProt ID</b>
	Human	<a href="#">30011</a> ;	<a href="#">Q96B97</a> ;
	Mouse		<a href="#">Q8R550</a> ;
	Rat		<a href="#">Q925Q9</a> ;
<b>Cellular Localization</b>	Cytoplasm . Cytoplasm, cytoskeleton. Cytoplasmic vesicle membrane; Peripheral membrane protein. Cell junction, synapse, synaptosome. Cell junction, focal adhesion . Localized in endocytic vesicles containing clustered receptors. Colocalizes with ASAP1 in vesicular structures. Colocalized with actin microfilaments and focal adhesions (By similarity). Colocalized with MAGI2 in synaptosomes. Translocation to EGFR containing vesicles upon EGF stimulation is inhibited in the presence of SH3KBP1 (By similarity). Colocalizes with ZFP36 in the cytoplasm (PubMed:20221403). .		
<b>Tissue specificity</b>	Ubiquitously expressed. Also expressed in some cancer cell lines.		
<b>Function</b>	<p>Domain:The SH3 domains mediate interaction with SHKBP1.,Function:Adapter protein involved in regulating diverse signal transduction pathways. Involved in the regulation of endocytosis and lysosomal degradation of ligand-induced receptor tyrosine kinases, including EGFR and MET/hepatocyte growth factor receptor, through a association with CBL and endophilins. The association with CBL, and thus the receptor internalization, may inhibited by an interaction with PDCD6IP and/or SPRY2. Involved in regulation of ligand-dependent endocytosis of the IgE receptor. Attenuates phosphatidylinositol 3-kinase activity by interaction with its regulatory subunit (By similarity). May be involved in regulation of cell adhesion; promotes the interaction between TTK2B and PDCD6IP. May be involved in the regulation of cellular stress response via the MAPK pathways through its interaction with MAP3K4. Is involved in modulation of tumor necrosis factor mediated apoptosis.,PTM:Monoubiquitinated by CBL and CBLB after EGF stimulation; probably on its C-terminus.,sequence Caution:Contaminating sequence. Potential poly-A sequence.,similarity:Contains 1 SH3 domain.,similarity:Contains 2 SH3 domains.,similarity:Contains 3 SH3 domains.,subcellular location:Localized in endocytic vesicles containing clustered receptors. Colocalizes with ASAP1 in vesicular structures. Colocalized with actin microfilaments and focal adhesions (By similarity). Colocalized with MAGI2 in synaptosomes.,subunit:Can self-associate and form homotetramers. Interacts with CD2, F-actin capping protein, PIK3R3, GRB2, EGFR, MET, BLNK, MAP3K4, PDCD6IP, SPRY2, ARHGAP17, ARHGAP27, MAGI2, CRK, BCAR1, SOS1, ASAP1, ARAP3, HIP1R, SYNJ2, INPP5D and STAP1. Interacts with CBL and CBLB, but does not interact with CBLC. Two molecules of SH3KBP1 seem to bind through their respective SH3 1 domain to one molecule of CBLB. The interaction with CBL or CBLB and EGFR is increased upon EGF stimulation. The interaction with CBL is attenuated by PDCD6IP. Interacts through its proline-rich region with the SH3 domain of endophilins SH3GL1, SH3GL2 and SH3GL3. The SH3KBP1-endophilin complex seems to associate with a complex containing the phosphorylated receptor (EGFR or MET) and phosphorylated CBL. Probably associates with ASAP1 and phosphorylated EGFR. Probably part of a complex consisting of at least SH3KBP1, ASAP1 and ARAP3. Interacts with focal adhesion kinases PTK2 AND PTK2B, probably as a dimer. Interacts with DAB2 and probably associates with chathrin through its interaction with DAB2. Part of a complex consisting of SH3KBP1, DAB2, and clathrin heavy chain. DAB2 and clathrin dissociate from SH3KBP1 following growth factor treatment, enabling interaction with CBL. Interacts with DDN and probably associates with MAGI2 through its interaction with DDN. Interacts with the SH3 domains of SRC tyrosine-protein kinases SRC, LCK, LYN, FGR, FYN and HCK. Interacts with TRADD, BIRC2, TRAF1, TRAF2 and TNFR1, and the association with a TNFR1-associated complex upon stimulation with TNF-alpha seems to be mediated by SRC. Probably interacts with SH3KBP1.,tissue specificity:Ubiquitously expressed. Also expressed in some cancer cell lines.,</p>		

---

## | Validation Data

## | Contact information

Orders: order@immunoway.com  
Support: tech@immunoway.com  
Telephone: 877-594-3616 (Toll Free), 408-747-0185  
Website: <http://www.immunoway.com>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code  
to access additional  
product information:  
**SH3K1 Rabbit pAb**

---

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)