

## JIP4 Rabbit pAb

CatalogNo: YN1592

### Key Features

**Host Species**

- Rabbit

**Reactivity**

- Human, Mouse

**Applications**

- WB, ELISA

**MW**

- 145kD (Observed)

**Isotype**

- IgG

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Recommended Dilution Ratios

**WB 1:500-2000****ELISA 1:5000-20000**

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** Synthesized peptide derived from human protein . at AA range: 590-670**Specificity** JIP4 Polyclonal Antibody detects endogenous levels of protein.

### Target Information

**Gene name** SPAG9 HSS KIAA0516 MAPK8IP4 SYD1 HLC6

**Protein Name** C-Jun-amino-terminal kinase-interacting protein 4 (JIP-4) (JNK-interacting protein 4) (Cancer/testis antigen 89) (CT89) (Human lung cancer oncogene 6 protein) (HLC-6) (JNK-associated leucine-zipper protein) (JLP) (Mitogen-activated protein kinase 8-interacting protein 4) (Proliferation-inducing protein 6) (Protein highly expressed in testis) (PHET) (Sperm surface protein) (Sperm-associated antigen 9) (Sperm-specific protein) (Sunday driver 1)

Organism	Gene ID	UniProt ID
Human	<a href="#">9043;</a>	<a href="#">O60271;</a>
Mouse		<a href="#">Q58A65;</a>

**Cellular Localization** Cytoplasm . Cytoplasm, perinuclear region . Lysosome membrane . Perinuclear distribution in response to stress signals such as UV radiation. . ; [Isoform 5]: Cytoplasmic vesicle, secretory vesicle, acrosome . Associated with the plasma membrane of the acrosomal compartment and also localizes in the acrosome matrix. .

**Tissue specificity** Expressed only in testis on the round spermatids of stage I, II and II. Absent in spermatogonia and spermatocyte. ; [Isoform 4]: Expressed in testis and in acute myeloid leukemia (AML) patients. ; [Isoform 3]: Expressed in testis.

**Function** Function:The JNK-interacting protein (JIP) group of scaffold proteins selectively mediates JNK signaling by aggregating specific components of the MAPK cascade to form a functional JNK signaling module. Isoform 5 may play a role in spermatozoa-egg-interaction.,induction:Isoform 3 is increased in systemic sclerosis fibroblasts.,PTM:Phosphorylated by MAPK8 and MAPK14 (By similarity). Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the JIP scaffold family.,subcellular location:Associated with the plasma membrane of the acrosomal compartment and also localizes in the acrosome matrix.,subcellular location:Perinuclear distribution in response to stress signals such as UV radiation.,subunit:Homo-oligomer. Interacts with MAX, MAPK8, MAPK9, MAPK10, MAPK14, MAP3K3, MYC, KNS2 and MAP2K4. Interaction with KNS2 is important in the formation of ternary complex with MAPK8. Interacts with NFKB1.,tissue specificity:Isoform 5 is expressed only in testis on the round spermatids of stage I, II and II. Isoform 5 is absent in spermatogonia and spermatocyte. Isoform 3 is expressed in testis. Isoform 4 is expressed in testis and in acute myeloid leukemia (AML) patients.,

---

## | Validation Data

## | Contact information

Orders: [order@immunoway.com](mailto:order@immunoway.com)  
Support: [tech@immunoway.com](mailto:tech@immunoway.com)  
Telephone: 877-594-3616 (Toll Free), 408-747-0185  
Website: <http://www.immunoway.com>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**JIP4 Rabbit pAb**

