Applications

WB,ELISA



NEB2 Rabbit pAb

CatalogNo: YN1686

Key Features

Host Species Reactivity

Rabbit
Human, Mouse, Rat

MW Isotype • 89kD (Observed) • IgG

Recommended Dilution Ratios

WB 1:500-2000 ELISA 1:5000-20000

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol,0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from human protein . at AA range: 660-740

Specificity NEB2 Polyclonal Antibody detects endogenous levels of protein.

| Target Information

Gene name PPP1R9B PPP1R6

Protein Name

Neurabin-2 (Neurabin-II) (Protein phosphatase 1 regulatory subunit 9B) (Spinophilin)

Organism	Gene ID	UniProt ID
Human	<u>84687</u> ;	Q96SB3;
Mouse		Q6R891;
Rat		<u>035274;</u>

Cellular Localization

Cytoplasm, cytoskeleton . Nucleus . Cell projection, dendritic spine . Cell junction, synapse, postsynaptic density . Cell junction, synapse. Cell junction, adherens junction . Cytoplasm. Cell membrane. Cell projection, lamellipodium. Cell projection, filopodium. Cell projection, ruffle membrane. Enriched at synapse and cadherin-based cell-cell adhesion sites. In neurons, both cytosolic and membrane-associated, and highly enriched in the postsynaptic density apposed to exitatory synapses. Colocalizes with PPP1R2 at actin-rich adherens junctions in epithelial cells and in dendritic spines (By similarity). Accumulates in the lamellipodium, filopodium and ruffle membrane in response to hepatocyte growth factor (HGF) treatment.

Tissue specificity Amygdala, Brain, Skin,

Function

Function: Seems to act as a scaffold protein in multiple signaling pathways. Modulates excitatory synaptic transmission and dendritic spine morphology. Binds to actin filaments (F-actin) and shows cross-linking activity. Binds along the sides of the F-actin. May play an important role in linking the actin cytoskeleton to the plasma membrane at the synaptic junction. Believed to target protein phosphatase 1/PP1 to dendritic spines, which are rich in F-actin, and regulates its specificity toward ion channels and other substrates, such as AMPA-type and NMDA-type glutamate receptors. Plays a role in regulation of G-protein coupled receptor signaling, including dopamine D2 receptors and alpha-adrenergic receptors. May establish a signaling complex for dopaminergic neurotransmission through D2 receptors by linking receptors downstream signaling molecules and the actin cytoskeleton. Binds to ADRA1B and RGS2 and mediates regulation of ADRA1B signaling. May confer to Rac signaling specificity by binding to both, RacGEFs and Rac effector proteins. Probably regulates p70 S6 kinase activity by forming a complex with TIAM1..PTM:Stimulation of D1 (but not D2) dopamine receptors induces Ser-94 phosphorylation. Dephosphorylation of Ser-94 is mediated mainly by PP1 and to a lesser extent by PP2A. Phosphorylation of spinophilin disrupts its association with F-actin, but does not affect its binding to PP1., similarity: Contains 1 PDZ (DHR) domain., subcellular location:Enriched at synapse and cadherin-based cell-cell adhesion sites. In neurons, both cytosolic and membrane-associated, and highly enriched in the post-synaptic density apposed to exitatory synapses. Colocalizes with PPP1R2 at actin-rich adherens junctions in epithelial cells and in dendritic spines., subunit: Possibly exists as an homodimer, homotrimer or an homotetramer. Interacts with F-actin, PPP1CA, neurabin-1, TGN38 and D(2) dopamine receptor. Interacts with RGS1, RGS2, RGS4, RGS19 and ADRA1B, ADRA2A, ADRA2B, ADRA2C, CDKN2A, PPP1R2, RASGFR1 and TIAM1.,

Validation Data

Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **NEB2 Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents