

NOX4 Rabbit pAb

CatalogNo: YN2975

Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB, ELISA

MW

- 63kD (Observed)

Isotype

- IgG

Recommended Dilution Ratios

WB 1:500-2000

ELISA 1:5000-20000

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen Synthesized peptide derived from part region of human protein AA range:520-578

Specificity NOX4 Polyclonal Antibody detects endogenous levels of protein.

Target Information

Gene name NOX4 RENOX

Protein Name	NADPH oxidase 4 (Kidney oxidase-1) (KOX-1) (Kidney superoxide-producing NADPH oxidase) (Renal NAD(P)H-oxidase)		
	Organism	Gene ID	UniProt ID
	Human	50507 ;	Q9NPH5 ;
	Mouse		Q9JHI8 ;
	Rat		Q924V1 ;
Cellular Localization	Endoplasmic reticulum membrane ; Multi-pass membrane protein. Cell membrane ; Multi-pass membrane protein . Cell junction, focal adhesion . May localize to plasma membrane and focal adhesions. According to PubMed:15927447, may also localize to the nucleus.; [Isoform 4]: Nucleus . Nucleus, nucleolus .		
Tissue specificity	Expressed by distal tubular cells in kidney cortex and in endothelial cells (at protein level). Widely expressed. Strongly expressed in kidney and to a lower extent in heart, adipocytes, hepatoma, endothelial cells, skeletal muscle, brain, several brain tumor cell lines and airway epithelial cells.		
Function	developmental stage:Expressed in fetal kidney and fetal liver.,enzyme regulation:Inhibited by plumbagin (By similarity). Activated by phorbol 12-myristate 13-acetate (PMA). Activated by insulin. Inhibited by diphenylene iodonium.,Function:Constitutive NADPH oxidase which generates superoxide intracellularly upon formation of a complex with CYBA/p22phox. Regulates signaling cascades probably through phosphatases inhibition. May function as an oxygen sensor regulating the KCNK3/TASK-1 potassium channel and HIF1A activity. May regulate insulin signaling cascade. May play a role in apoptosis, bone resorption and lipopolysaccharide-mediated activation of NFkB. Isoform 3 is not functional. Isoform 4 displays an increased activity while isoform 5 and isoform 6 display reduced activity. May produce superoxide in the nucleus and play a role in regulating gene expression upon cell stimulation.,induction:By 7-ketocholesterol (at protein level).,PTM:Isoform 3 and isoform 4 are N-glycosylated. Isoform 4 glycosylation is required for its proper function.,similarity:Contains 1 FAD-binding FR-type domain.,similarity:Contains 1 ferric oxidoreductase domain.,subcellular location:May localize to plasma membrane and focal adhesions. May also localize to the nucleus (PubMed:15927447).,subunit:Interacts with protein disulfide isomerase (By similarity). Interacts with, relocalizes and stabilizes CYBA/p22phox. Interacts with TLR4.,tissue specificity:Expressed by distal tubular cells in kidney cortex and in endothelial cells (at protein level). Widely expressed. Strongly expressed in kidney and to a lower extent in heart, adipocytes, hepatoma, endothelial cells, skeletal muscle, brain, several brain tumor cell lines and airway epithelial cells.,		

| Validation Data

| Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code
to access additional
product information:
NOX4 Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)