**Applications** 

WB



## **ACY2 Rabbit pAb**

CatalogNo: YN3763

# **| Key Features**

Host Species Reactivity

Rabbit
 Human, Mouse, Rat

MW Isotype
• 34kD (Calculated) • IgG

### **Recommended Dilution Ratios**

WB 1:500-2000

### Storage

Storage\* -15°C to -25°C/1 year(Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### **Basic Information**

**Clonality** Polyclonal

## Immunogen Information

**Immunogen** Synthesized peptide derived from human ACY2 AA range: 67-117

**Specificity** This antibody detects endogenous levels of ACY2 at Human/Mouse/Rat

### | Target Information

**Gene name** ASPA ACY2 ASP

#### **Protein Name**

ACY2

Organism	Gene ID	UniProt ID
Human	<u>443</u> ;	<u>P45381;</u>
Mouse	<u>11484;</u>	<u>Q8R3P0</u> ;
Rat	<u>79251;</u>	<u>Q9R1T5;</u>

#### Cellular Localization


Cytoplasm. Nucleus.

**Tissue specificity** Brain white matter, skeletal muscle, kidney, adrenal glands, lung and liver.

#### **Function**

Catalytic activity:N-acyl-L-aspartate + H(2)O = a carboxylate + L-aspartate.,cofactor:Binds 1 zinc ion per subunit.,Disease:Defects in ASPA are the cause of Canavan disease (CAND) [MIM:271900]; also known as spongy degeneration of the brain. CAND is a rare neurodegenerative condition of infancy or childhood characterized by white matter vacuolization and demeylination that gives rise to a spongy appearance. The clinical features are onset in early infancy, atonia of neck muscles, hypotonia, hyperextension of legs and flexion of arms, blindness, severe mental defect, megalocephaly, and death by 18 months on the average.,Function:Catalyzes the deacetylation of N-acetylaspartic acid (NAA) to produce acetate and L-aspartate. NAA occurs in high concentration in brain and its hydrolysis NAA plays a significant part in the maintenance of intact white matter. In other tissues it act as a scavenger of NAA from body fluids.,similarity:Belongs to the aspA/astE family. Aspartoacylase subfamily.,subunit:Homodimer.,tissue specificity:Brain white matter, skeletal muscle, kidney, adrenal glands, lung and liver.,

### **| Validation Data**



Western blot analysis of lysates from 3T3 cells, primary antibody was diluted at 1:1000, 4°over night

### **Contact information**

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **ACY2 Rabbit pAb** 

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents