

GLSK Rabbit pAb

CatalogNo: YN4051

| Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB

MW

- 74kD (Calculated)

Isotype

- IgG

| Recommended Dilution Ratios

WB 1:500-2000

| Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

| Basic Information

Clonality Polyclonal

| Immunogen Information

Immunogen Synthesized peptide derived from human GLSK AA range: 162-212

Specificity This antibody detects endogenous levels of GLSK at Human/Mouse/Rat

| Target Information

Gene name GLS GLS1 KIAA0838

Protein Name GLSK

Organism	Gene ID	UniProt ID
Human	2744;	O94925;
Mouse	14660;	D3Z7P3;
Rat	24398;	P13264;

**Cellular
Localization**

[Isoform 1]: Mitochondrion . Cytoplasm, cytosol . The 74-kDa cytosolic precursor is translocated into the mitochondria and processed via a 72-kDa intermediate to yield the mature 68- and 65-kDa subunits. .; [Isoform 3]: Mitochondrion .; [Glutaminase kidney isoform, mitochondrial 68 kDa chain]: Mitochondrion matrix . Produced by the proteolytic processing of the 74-kDa cytosolic precursor. .; [Glutaminase kidney isoform, mitochondrial 65 kDa chain]: Mitochondrion matrix . Produced by the proteolytic processing of the 74-kDa cytosolic precursor. .

Tissue specificity

Isoform 1 and isoform 3 are detected in brain cortex. Isoform 3 is highly expressed in astrocytoma, ganglioglioma and ependymoma. Isoform 1 is highly expressed in brain and kidney, but not detected in liver. Isoform 3 is highly expressed in heart and pancreas, detected at lower levels in placenta, lung, pancreas and kidney, but is not detected in liver. Isoform 2 is expressed in cardiac and skeletal muscle.

Function

Catalytic activity:L-glutamine + H(2)O = L-glutamate + NH(3).,Function:Catalyzes the first reaction in the primary pathway for the renal catabolism of glutamine.,similarity:Belongs to the glutaminase family.,similarity:Contains 1 ANK repeat.,tissue specificity:KGA is expressed predominantly in brain and kidney but not in liver, GAC is expressed principally in cardiac muscle and pancreas but not in liver or brain, and GAM is expressed solely in cardiac and skeletal muscle.,

| Validation Data

| Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com
Telephone: 877-594-3616 (Toll Free), 408-747-0185
Website: <http://www.immunoway.com>
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
GLSK Rabbit pAb