

MAP1S Rabbit pAb

CatalogNo: YN6145

Key Features

Host Species
• Rabbit
MW

116kD (Calculated)

Reactivity
• Human,Mouse,Rat

Applications
• WB

Isotype • IgG

Recommended Dilution Ratios

WB 1:500-2000

Storage

Storage*	-15°C to -25°C/1 year(Do not lower than -25°C)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen	Synthesized peptide derived from human MAP1S
Specificity	This antibody detects endogenous levels of MAP1S at Human, Mouse,Rat

Target Information

Gene name MAP1S BPY2IP1 C19orf5 MAP8 VCY2IP1

Microtubule-associated protein 1S (MAP-1S) (BPY2-interacting protein 1) (Microtubuleassociated protein 8) (Variable charge Y chromosome 2-interacting protein 1) (VCY2interacting protein 1) (VCY2IP-1) [Cleaved into: MAP1S heavy chain; MAP1S light chain]

Organism	Gene ID	UniProt ID
Human	<u>55201;</u>	<u>Q66K74;</u>
Mouse	<u>270058;</u>	<u>Q8C052;</u>
Rat	<u>290640;</u>	<u>P0C5W1;</u>

- **Cellular** Localization Nucleus. Cytoplasm, cytosol. Cytoplasm, cytoskeleton . Cytoplasm, cytoskeleton, spindle . Detected in filopodia-like protrusions and synapses (By similarity). Detected in perinuclear punctate network corresponding to mitochondrial aggregates and in the nucleus in cells exhibiting apoptosis. Associated specifically with microtubules stabilized by paclitaxel and colocalizes with RASSF1 isoform A. In interphase cells, shows a diffuse cytoplasmic staining with partial localization to the microtubules. During the different stages of mitosis detected at the spindle microtubules. .
- **Tissue specificity** Expressed in neurons (at protein level). Expressed in spermatocytes, spermatids and spermatozoa. Expressed in the cerebral cortex. Highly expressed in testis. Moderately expressed in the brain, colon, heart, kidney, liver, lung, placenta, small intestine, spleen and stomach. Weakly expressed in muscle.
- **Function** Microtubule-associated protein that mediates aggregation of mitochondria resulting in cell death and genomic destruction (MAGD). Plays a role in anchoring the microtubule organizing center to the centrosomes. Binds to DNA. Plays a role in apoptosis. Involved in the formation of microtubule bundles (By similarity).

Validation Data

Contact information

Orders:	order@immunoway.com
Support:	tech@immunoway.com
Telephone:	877-594-3616 (Toll Free), 408-747-0185
Website:	http://www.immunoway.com
Address:	2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information: **MAP1S Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents