

RFFL Rabbit pAb

CatalogNo: YN8265

| Key Features

Host Species

- Rabbit

Reactivity

- Human, Mouse, Rat

Applications

- WB

MW

- 40kD (Calculated)

Isotype

- IgG

| Recommended Dilution Ratios

WB 1:500-2000

| Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

| Basic Information

Clonality Polyclonal

| Immunogen Information

Immunogen Synthesized peptide derived from human RFFL

Specificity This antibody detects endogenous levels of RFFL at Human, Mouse, Rat

| Target Information

Gene name RFFL RNF189 RNF34L

Protein Name	E3 ubiquitin-protein ligase rififylin (Caspase regulator CARP2) (Caspases-8 and -10-associated RING finger protein 2) (CARP-2) (FYVE-RING finger protein Sakura) (Fring) (RING finger and FYVE-like domain-containing protein 1) (RING finger protein 189) (RING finger protein 34-like)		
	Organism	Gene ID	UniProt ID
	Human	117584 ;	Q8WZ73 ;
	Mouse	67338 ;	Q6ZQM0 ;
	Rat	282844 ;	Q8CIN9 ;
Cellular Localization	Cytoplasm, cytosol. Cell membrane; Peripheral membrane protein. Recycling endosome membrane ; Peripheral membrane protein . The FYVE-type zinc finger may mediate phosphatidylinositol phosphate-binding and control subcellular localization.		
Tissue specificity	Ubiquitous. Detected in spleen, thymus, prostate, testis, ovary, small intestine, colon and peripheral blood leukocytes.		
Function	E3 ubiquitin-protein ligase that regulates several biological processes through the ubiquitin-mediated proteasomal degradation of various target proteins. Mediates 'Lys-48'-linked polyubiquitination of PRR5L and its subsequent proteasomal degradation thereby indirectly regulating cell migration through the mTORC2 complex. Ubiquitinates the caspases CASP8 and CASP10, promoting their proteasomal degradation, to negatively regulate cell death downstream of death domain receptors in the extrinsic pathway of apoptosis. Negatively regulates the tumor necrosis factor-mediated signaling pathway through targeting of RIPK1 to ubiquitin-mediated proteasomal degradation. Negatively regulates p53/TP53 through its direct ubiquitination and targeting to proteasomal degradation. Indirectly, may also negatively regulate p53/TP53 through ubiquitination and degradation of SFN. May also play a role in endocytic recycling.		

| Validation Data

| Contact information

Orders: order@immunoway.com
 Support: tech@immunoway.com
 Telephone: 877-594-3616 (Toll Free), 408-747-0185
 Website: <http://www.immunoway.com>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
RFFL Rabbit pAb