Applications

WB,IHC,IF,ELISA



Cytokeratin 8 (Phospho Ser73) Rabbit pAb

CatalogNo: YP0082 Orthogonal Validated 👀

Key Features

Host Species Reactivity

RabbitHuman, Mouse

MW Isotype
• 55kD (Observed) • IgG

Recommended Dilution Ratios

WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:40000 IF 1:50-200

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Basic Information

Clonality Polyclonal

Immunogen Information

Immunogen The antiserum was produced against synthesized peptide derived from human Keratin 8

around the phosphorylation site of Ser73. AA range:41-90

Specificity

Phospho-Cytokeratin 8 (S73) Polyclonal Antibody detects endogenous levels of Cytokeratin 8 protein only when phosphorylated at S73. The name of modified sites may be influenced by many factors, such as species (the modified site was not originally found in human samples) and the change of protein sequence (the previous protein sequence is incomplete, and the protein sequence may be prolonged with the development of protein sequencing technology). When naming, we will use the "numbers" in historical reference to keep the sites consistent with the reports. The antibody binds to the following modification sequence (lowercase letters are modification sites):LLsPL

Target Information

Gene name

KRT8 CYK8

Protein Name

CARD2;CK 8;CK-8;CK8;CYK8;CYKER;Cytokeratin endo

A;Cytokeratin-8;DreK8;EndoA;K2C8;K2C8 HUMAN;K8;Keratin 8;Keratin type II cytoskeletal

8; Keratin, type II cytoskeletal 8; Keratin-8; KO; Krt

2.8;KRT8;MGC118110;MGC174782;MGC53564;MGC85764;sb:cb186;Type-II keratin Kb8

Organism	Gene ID	UniProt ID
Human	<u>3856;</u>	<u>P05787;</u>
Mouse	<u>16691;</u>	<u>P11679;</u>
Rat	<u>25626;</u>	<u>Q10758</u> ;

Cellular Localization

Cytoplasm . Nucleus, nucleoplasm . Nucleus matrix .


Tissue specificity Observed in muscle fibers accumulating in the costameres of myoplasm at the sarcolemma membrane in structures that contain dystrophin and spectrin. Expressed in gingival mucosa and hard palate of the oral cavity.

Function


Disease:Defects in KRT8 are a cause of cryptogenic cirrhosis [MIM:215600]., Function: Together with KRT19, helps to link the contractile apparatus to dystrophin at the costameres of striated muscle., miscellaneous: There are two types of cytoskeletal and microfibrillar keratin: I (acidic; 40-55 kDa) and II (neutral to basic; 56-70 kDa).,PTM:O-glycosylated at multiple sites; glycans consist of single N-acetylglucosamine residues., PTM: Phosphorylation on serine residues is enhanced during EGF stimulation and mitosis. Ser-74 phosphorylation plays an important role in keratin filament reorganization., similarity: Belongs to the intermediate filament

family., subunit: Heterotetramer of two type I and two type II keratins. keratin-8 associates with keratin-18. Associates with KRT20. Interacts with HCV core protein and PNN. When associated with KRT19, interacts with DMD. Interacts with TCHP., tissue specificity: Observed in muscle fibers accumulating in the costameres of myoplasm at the sarcolemma membrane in structures that contain dystrophin and spectrin. Expressed in gingival mucosa


and hard palate of the oral cavity.,




Western Blot analysis of 293 cells using Phospho-Cytokeratin 8 (S73) Polyclonal Antibody diluted at 1:500



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Keratin 8 (Phospho-Ser73) Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using Keratin 8 (Phospho-Ser73) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from 293 cells treated with Etoposide 25uM 60', using Keratin 8 (Phospho-Ser73) Antibody. The lane on the right is blocked with the phospho peptide.

| Contact information

Orders: order@immunoway.com Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

Cytokeratin 8
(Phospho Ser73)

Rabbit pAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Antibody | ELISA Kits | Protein | Reagents